arXiv:1411.2556v1 [quant-ph] 7 Nov 2014

On geometry influence on the behavior of a quantum mechanical scalar particle with intrinsic structure in external magnetic and electric fields

November 11, 2014

O.V. Veko[1], K.V. Kazmerchuk[2], E.M. Ovsiyuk[3], V.V. Kisel[4], V.M. Red'kov[5]

Abstract

Relativistic theory of the Cox's scalar not point-like particle with intrinsic structure is developed on the background of arbitrary curved space-time. It is shown that in the most general form, the extended Proca-like tensor first order system of equations contains non minimal interaction terms through electromagnetic tensor $F_{\alpha\beta}$ and Ricci tensor $R_{\alpha\beta}$.

In relativistic Cox's theory, the limiting procedure to non-relativistic approximation is performed in a special class of curved space-time models. This theory is specified in simple geometrical backgrounds: Euclid's, Lobachevsky's, and Riemann's. Wave equation for the Cox's particle is solved exactly in presence of external uniform magnetic and electric fields in the case of Minkowski space. Non-trivial additional structure of the particle modifies the frequency of a quantum oscillator arising effectively in presence if external magnetic field. Extension of these problems to the case of the hyperbolic Lobachevsky space is examined. In presence of the magnetic field, the quantum problem in radial variable has been solved exactly; the quantum motion in z-direction is described by 1-dimensional Schrödinger-like equation in an effective potential which turns out to be too difficult for analytical treatment. In the presence of electric field, the situation is similar. The same analysis has been performed for spherical Riemann space model.

PACS numbers: 02.30.Gp, 02.40.Ky, 03.65Ge, 04.62.+v
MSC 2010: 33E30, 34B30
KeywordsIntrinsic structure, scalar particle, curved space-time, generalized Schrödinger equation, magnetic field, electric field, Minkowski, Lobachevsky, Riemann space models

1 Scalar Cox's particle with intrinsic structure

In 1982 W. Cox [1] proposed a special wave equation for a scalar particle with a larger set of tensor components than the usual Proca's approach includes: he used the set of a scalar, 4-vector, antisymmetric and (irreducible) symmetric tensor, thus starting with the 20-component wave function (see Section **13**).

[1]Kalinkovichi Gymnasium, Belarus,vekoolga@mail.ru
[2]Mosyr State Pedagogical University, Belarus, kristinash2@mail.ru
[3]Mosyr State Pedagogical University, Belarus, e.ovsiyuk@mail.ru
[4]Belarusian State University of Informatics and Radioelectronics
[5]B.I. Stepanov Institute of Physics, NAS of Belarus, redkov@dragon.bas-net.by

First, let us consider the system of Cox's equations [1] in the Minkowski space. We will use a Proca-like generalized system obtained after elimination from the initial system of Cox's equations two second-rank tensors (see Section **13**):

$$\left(\frac{i}{\sqrt{-g}}\,\frac{\partial}{\partial x^{\alpha}}\sqrt{-g}-\frac{e}{c\hbar}\,A_{\alpha}\right)\,g^{\alpha\beta}\Phi_{\beta}=\frac{Mc}{\hbar}\Phi\;,\quad K_{\rho}{}^{\alpha}\left(i\,\partial_{\alpha}-\frac{e}{c\hbar}A_{\alpha}\right)\Phi=\frac{Mc}{\hbar}\Phi_{\rho}\;. \tag{1.1}$$

$K_{\rho}{}^{\alpha}$ is a tensor inverse to $\Lambda_{\sigma}{}^{\alpha}=Mc\delta_{\sigma}{}^{\alpha}+\lambda F_{\sigma}{}^{\alpha}$ (λ stands for additional parameter responsible for non-trivia intrinsic structure of a scalar particle in Cox's approach):

$$K_{\rho}{}^{\alpha}=\lambda_1\,\delta_{\alpha}{}^{\beta}+\lambda_2\,F_{\alpha}{}^{\beta}+\lambda_3\,F_{\alpha}{}^{\rho}F_{\rho}{}^{\beta}+\lambda_4\,F_{\alpha}{}^{\rho}F_{\rho}{}^{\sigma}F_{\sigma}{}^{\beta}\;, \tag{1.2}$$

λ_i are expressed through electromagnetic invariants (for more technical details see Section **13**).

In geometrical models with metrics of special type $dS^2=c^2dt^2+g_{kl}(x)\,dx^k dx^l$, one cap perform non-relativistic approximation and derive extended Schrödinger type equation (see Section **13**):

$$D_t\,\Psi=\frac{1}{2M}\,\overset{\circ}{D}_k\,(-g^{kj})\left(K_j{}^{l}D_l+McK_j{}^{0}\right)\Psi-\frac{1}{2}\left((K_0{}^{0}-1)Mc^2+K_0{}^{j}cD_j\right)\Psi\;, \tag{1.3}$$

where the notation is used

$$i\hbar\,\partial_t-eA_0=D_t\;,\quad ic\hbar\,\partial_k-eA_k=cD_k\;,\quad \frac{ic\hbar}{\sqrt{-g}}\,\frac{\partial}{\partial x^k}\sqrt{-g}-e\,A_k=c\,\overset{\circ}{D}_k\;,$$

It is a generalized Schrödinger equation for the particle with intrinsic structure.

In presence of a pure magnetic field, the above equation (1.3) takes a more simple form

$$D_t\,\Psi=-\frac{1}{2M}\,\overset{\circ}{D}_k\,g^{kj}(x)\,\overset{*}{D}_j\,\Psi\;, \tag{1.4}$$

where the notation is used (let $\Gamma=\lambda/mc$)

$$\begin{aligned}
\overset{*}{D}_1&=K_1{}^{l}D_l=\frac{1}{1+\Gamma^2B_iB^i}\left[D_1+\Gamma(B_2D^3-B_3D^2)+\Gamma^2B^1\,(B_iD_i)\right],\\
\overset{*}{D}_2&=K_2{}^{l}D_l=\frac{1}{1+\Gamma^2B_iB^i}\left[D_2+\Gamma(B_3D^1-B_1D^3)+\Gamma^2B^2\,(B_iD_i)\right],\\
\overset{*}{D}_3&=K_3{}^{l}D_l=\frac{1}{1+\Gamma^2B_iB^i}\left[D_3+\Gamma(B_1D^2-B_2D^1)+\Gamma^2B^3\,(B_iD_i)\right],\\
&g^{22}g^{33}B_1=B^1\;,\quad g^{33}g^{11}B_2=B^2\;,\quad g^{11}g^{22}B_3=B^3\;,\quad F_{ij}=\epsilon_{ijk}B_k\;.
\end{aligned}$$

In presence of a pure electric field, the above equation (1.3) takes the form

$$\left(D_t-c\,\frac{\Gamma^2E_iE^iMc+\Gamma E^jD_j}{2(1+\Gamma^2E_iE^i)}\right)\Psi=\frac{1}{2M}\,\overset{\circ}{D}_k\,(-g^{kj})\left(D_j+\frac{\Gamma^2E_j(E^iD_i)+Mc\Gamma E_j}{1+\Gamma^2E_iE^i}\right)\Psi \tag{1.5}$$

where

$$g^{11}E_1=E^1,\qquad g^{22}E_2=E^2,\qquad g^{33}E_3=E^3,\qquad E_i=F_{0i}\;.$$

2 Cox's particle in the magnetic field, Minkowski space

Let the homogeneous magnetic field $\vec{A} = \frac{1}{2}\vec{x} \times \vec{B}$ be directed along the axis z:

$$(A_j) = \vec{A} = \frac{1}{2}\left(x^2 B,\ -x^1 B,\ 0\right)\ .$$

Recalculating the potential to cylindrical coordinates by the formulas

$$A_{j'} = \frac{\partial x^j}{\partial x^{j'}} A_j,\ x^j = (x, y, z),\ xj' = (r, \phi, z)\ , \quad x^1 = r\cos\phi,\ x^2 = r\sin\phi,\ x^3 = x^{3'} = z\ ,$$

we obtain

$$A_r = 0,\ A_\phi = -\frac{Br^2}{2},\ A_z = 0,\ F_{r\phi} = -Br\ . \tag{2.1}$$

The metric tensor in these coordinates and field variables are determined by

$$dS^2 = c^2dt^2 - dr^2 - r^2d\phi^2 - dz^2,\ \sqrt{-g} = r\ , \quad B_3 = -Br,\ B^3 = -Br^{-1},\ B_iB^i = B^2\,. \tag{2.2}$$

The Schrödinger equation for this case reads

$$D_t\Psi = \frac{1}{2M}\left(\overset{\circ}{D}_1\overset{*}{D}_1 + \overset{\circ}{D}_2\, r^{-2}\ \overset{*}{D}_2 + \overset{\circ}{D}_3\overset{*}{D}_3\right)\Psi,$$

where

$$D_1 = i\hbar\partial_r\ ,\ D_2 = i\hbar\partial_\phi + \frac{e}{c}\frac{Br^2}{2}\ ,\ D_3 = i\hbar\partial_z\ , \quad \overset{\circ}{D}_1 = i\hbar(\partial_r + \frac{1}{r}), \qquad \overset{\circ}{D}_2 = i\hbar\partial_\phi + \frac{e}{c}\frac{Br^2}{2}\ ,$$
$$\overset{\circ}{D}_3 = i\hbar\partial_z\ , \qquad \overset{*}{D}_1 = \frac{1}{1+\Gamma^2B^2}(D_1 - \Gamma B_3D^2) = \frac{1}{1+\Gamma^2B^2}\left(i\hbar\partial_r - \frac{\Gamma B}{r}(i\hbar\partial_\phi + \frac{e}{c}\frac{Br^2}{2})\right),$$

$$\overset{*}{D}_2 = \frac{1}{1+\Gamma^2B^2}(\ D_2 + \Gamma B_3D^1) = \frac{1}{1+\Gamma^2B^2}\left((i\hbar\partial_\phi + \frac{e}{c}\frac{Br^2}{2}) + i\hbar\Gamma Br\partial_r\right),$$

$$\overset{*}{D}_3 = \frac{(D_3 + \Gamma^2B^3\ B_3D_3)}{1+\Gamma^2B^2} = i\hbar\partial_z\ ;$$

below we will use the notation

$$\frac{eB}{2\hbar c} = b, \qquad \Gamma B = \gamma\ .$$

We compute

$$\frac{1}{2M}\ \overset{\circ}{D}_1\overset{*}{D}_1 = -\frac{\hbar^2}{2M(1+\gamma^2)}\left(\partial_r^2 + \frac{1}{r}\partial_r - \frac{\gamma}{r}\partial_r\partial_\phi + i\gamma br\partial_r + 2i\gamma b\right),$$
$$\frac{1}{2M}\ \overset{\circ}{D}_2\overset{*}{D}_2 = -\frac{\hbar^2}{2M(1+\gamma^2)}\left(\frac{1}{r^2}(\partial_\phi - ibr^2)^2 + \gamma(\partial_\phi - ibr^2)\ \frac{1}{r}\partial_r\right),$$
$$\frac{1}{2M}\ \overset{\circ}{D}_3\overset{*}{D}_3 = -\frac{\hbar^2}{2M(1+\gamma^2)}(1+\gamma^2)\partial_z^2\ . \tag{2.3}$$

After using the substitution for the wave function

$$\Psi = e^{-iEt/\hbar}e^{im\phi}e^{ikz}R(r), \quad \epsilon = \frac{2mE}{\hbar^2}(1+\gamma^2)$$

we get the radial Schrödinger equation

$$\left[\frac{d^2}{dr^2}+\frac{1}{r}\frac{d}{dr}+2i\gamma b+\epsilon-\frac{(m-br^2)^2}{r^2}-(1+\gamma^2)k^2\right]R=0. \tag{2.4}$$

By physical reasons parameter γ must be purely imaginary (see Section **13**): $\gamma=-i\eta$; so the radial equation reads

$$\left[\frac{d^2}{dr^2}+\frac{1}{r}\frac{d}{dr}+2\eta b+\epsilon-\frac{(m-br^2)^2}{r^2}-(1-\eta^2)k^2\right]R=0. \tag{2.5}$$

With the use of notation $\epsilon-(1-\eta^2)k^2+2\eta b=\epsilon'$, equation (2.5) can be written as

$$\left(\frac{d^2}{dr^2}+\frac{1}{r}\frac{d}{dr}-\frac{(m-br^2)^2}{r^2}+\epsilon'\right)=0\ , \tag{2.6}$$

which coincides with the equation arising in the problem of the usual particle in the magnetic field. Its solutions are known. We present here only an expression for the energy spectrum

$$\epsilon=4b(n+\frac{m+\mid m\mid+1}{2})+(1-\eta^2)k^2-2\eta b\ ; \tag{2.7}$$

from this after translating to ordinary units we obtain

$$E=\frac{p^2}{2M}+\frac{1}{1-\eta^2}\frac{eB}{Mc}\hbar\left(n+\frac{m+\mid m\mid+1}{2}\right)-\frac{\eta}{2}\frac{1}{1-\eta^2}\frac{eB}{Mc}\hbar\ . \tag{2.8}$$

With the use of notation $\eta=\Gamma B,\ \Gamma^*=\Gamma,\ \omega=\frac{eB}{Mc}$, the formula for the energy levels can be written as

$$E=\frac{p^2}{2M}+\frac{\omega\hbar}{1-(\Gamma B)^2}\left(n+\frac{m+\mid m\mid+1}{2}\right)-\frac{\omega\hbar}{1-(\Gamma B)^2}\frac{\Gamma B}{2}\ . \tag{2.9}$$

Thus, the intrinsic structure of the Cox's particle modifies the frequency of the quantum oscillator (in fact, this result was firstly produced in different formalism by Kisel [2]).

$$\omega\ \Longrightarrow\ \tilde{\omega}=\frac{\omega}{1-\Gamma^2B^2},\ \omega=\frac{eB}{Mc}\ . \tag{2.10}$$

3 Cox's particle in the magnetic field in the Lobachevsky space

In a special (cylindrical) coordinate system in the Lobachevsky space, analogue of the uniform magnetic field is determined by the relations (we use dimensionless coordinate r obtained by dividing on the curvature radius ρ):

$$dS^2=c^2dt^2-\mathrm{ch}^2z(dr^2+\mathrm{sh}^2rd\phi^2)+dz^2,$$
$$A_\phi=-B\rho^2(\mathrm{ch}\ r-1),\ F_{r\phi}=-B\rho\ \mathrm{sh}\ r\ ,$$

$$B_3=-B\rho\ \mathrm{sh}\ r\ ,\ B^3=-\frac{B}{\rho\ \mathrm{sh}\ r\ \mathrm{ch}^4z}\ ,\quad B_iB^i=B^2\mathrm{ch}^{-4}z\ . \tag{3.1}$$

The wave equation in this case reads

$$D_t\Psi = \frac{1}{2M\rho^2}\left(\overset{\circ}{D}_1\overset{*}{D}_1 + \overset{\circ}{D}_2\,\frac{1}{\mathrm{sh}^2 r}\,\overset{*}{D}_2 + \overset{\circ}{D}_3\overset{*}{D}_3\right)\Psi, \tag{3.2}$$

where

$$D_1 = i\hbar\partial_r, \quad D_3 = i\hbar\partial_z, \quad D_2 = i\hbar\partial_\phi + \frac{e}{c}B\rho^2(\mathrm{ch}\ r - 1),$$

$$\overset{\circ}{D}_1 = i\hbar(\partial_r + \frac{\mathrm{ch}\ r}{\mathrm{sh}\ r}), \quad \overset{\circ}{D}_2 = i\hbar\partial_\phi + \frac{e}{c}B\rho^2(\mathrm{ch}\ r - 1), \quad \overset{\circ}{D}_3 = i\hbar(\partial_z + 2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z}),$$

$$\overset{*}{D}_1 = \frac{1}{1+\Gamma^2B^2\mathrm{ch}^{-4}z}\left[i\hbar\partial_r - \frac{\Gamma B\mathrm{ch}^{-2}z}{\mathrm{sh}r}(i\hbar\partial_\phi + \frac{e}{c}B\rho^2(\mathrm{ch}\ r-1))\right],$$

$$\overset{*}{D}_2 = \frac{1}{1+\Gamma^2B^2\mathrm{ch}^{-4}z}\left[(i\hbar\partial_\phi + \frac{e}{c}B\rho^2(\mathrm{ch}\ r-1)) + i\hbar\Gamma B\mathrm{ch}^{-2}z\ \mathrm{sh}\ r\partial_r\right],$$

$$\overset{*}{D}_3 = \frac{(D_3 + \Gamma^2B^3\ B_3D_3)}{1+\Gamma^2B^2\mathrm{ch}^{-4}z} = i\hbar\partial_z.$$

Below the notation is used:

$$(eB\rho^2/\hbar c) = b, \qquad \Gamma B\mathrm{ch}^{-2}z = \gamma(z).$$

We compute

$$\begin{aligned}
\frac{1}{2M\rho^2}\ \overset{\circ}{D}_1\ g^{11}\ \overset{*}{D}_1 &= -\frac{\hbar^2\mathrm{ch}^{-2}z}{2M\rho^2(1+\gamma^2(z))} \\
\times\left(\partial_r^2 + (\frac{\mathrm{ch}\ r}{\mathrm{sh}\ r} + i\gamma(z)b\frac{\mathrm{ch}\ r - 1}{\mathrm{sh}\ r})\partial_r - \frac{\gamma(z)}{\mathrm{sh}\ r}\partial_r\partial_\phi + i\gamma(z)b\right), \\
\frac{1}{2M\rho^2}\ \overset{\circ}{D}_2\ g^{22}\ \overset{*}{D}_2 &= -\frac{\hbar^2\mathrm{ch}^{-2}z}{2M\rho^2(1+\gamma^2(z))} \\
\times\left[\frac{1}{\mathrm{sh}^2 r}[\partial_\phi - ib(\mathrm{ch}\ r-1)]^2 + \gamma(z)[\partial_\phi - ib(\mathrm{ch}\ r-1)]\ \frac{1}{\mathrm{sh}\ r}\partial_r\right], \\
\frac{1}{2M\rho^2}\ \overset{\circ}{D}_3\ g^{33}\ \overset{*}{D}_3 &= -\frac{\hbar^2}{2M\rho^2}(\partial_z + 2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z})\partial_z.
\end{aligned} \tag{3.3}$$

After using the substitution for the wave function:

$$\Psi = e^{-iEt/\hbar}e^{im\phi}Z(z)R(r)\ , \tag{3.4}$$

the Schrödinger equation (3.2) gives (the function $\gamma(z)$ must be imaginary, $i\gamma(z \Longrightarrow \gamma(z))$

$$\begin{aligned}
&\left[\frac{\mathrm{ch}^{-2}z}{1-\gamma^2(z)}\left(\partial_r^2 + \frac{\mathrm{ch}\ r}{\mathrm{sh}\ r}\partial_r - \frac{[m - b(\mathrm{ch}\ r-1)]^2}{\mathrm{sh}^2 r} + b\gamma(z)\right)\right. \\
&\left. +\epsilon + (\partial_z + 2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z})\partial_z\right]R(r)Z(z) = 0\ , \quad \epsilon = \frac{E}{\hbar^2/2M\rho^2}\ .
\end{aligned} \tag{3.5}$$

In this equation, the variables are separated:

$$\frac{1}{R}\left(\frac{d^2}{dr^2}+\frac{\mathrm{ch}\ r}{\mathrm{sh}\ r}\frac{d}{dr}-\frac{[m-b(\mathrm{ch}\ r-1)]^2}{\mathrm{sh}^2 r}\right)R$$
$$+\frac{1}{Z}(1-\gamma^2(z))\ \mathrm{ch}^2 z\left(\frac{b\,\gamma(z)\ \mathrm{ch}^{-2}z}{1-\gamma^2(z)}+\epsilon+(\frac{d}{dz}+2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z})\frac{d}{dz}\right)Z=0. \tag{3.6}$$

The radial equation for the function $R(r)$ reads

$$\left(\frac{d^2}{dr^2}+\frac{\mathrm{ch}\ r}{\mathrm{sh}\ r}\frac{d}{dr}-\frac{[m-b(\mathrm{ch}\ r-1)]^2}{\mathrm{sh}^2 r}+\Lambda\right)R=0\ ; \tag{3.7}$$

the equation for $Z(z)$ is (remember that $\gamma = B\Gamma$)

$$\left(\frac{d^2}{dz^2}+2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z}\frac{d}{dz}+\epsilon+\frac{b\gamma-\Lambda\mathrm{ch}^2 z}{\mathrm{ch}^4 z-\gamma^2}\right)Z=0.$$

4 Analysis of the equation in the variable z

In equation (3.8), let us eliminate the first derivative term:

$$Z=\frac{1}{\mathrm{ch}\ z}f(z),\quad U(z)=-\frac{b\gamma-\Lambda\ \mathrm{ch}^2 z}{\mathrm{ch}^4 z-\gamma^2},$$
$$\left(\frac{d^2}{dz^2}+\epsilon-1-U(z)\right)f(z)=0. \tag{4.1}$$

Eq. (4.1) can be viewed as the Schrödinger equation in the effective potential field $U(z)$. The corresponding effective force is

$$F_z=-\frac{dU}{dz}$$
$$=2\ \mathrm{ch}\ z\ \mathrm{sh}\ z\ \frac{\Lambda\ \mathrm{ch}^4 z-2b\gamma\ \mathrm{ch}^2 z+\gamma^2\Lambda}{(\mathrm{ch}^4 z-\gamma^2)^2}. \tag{4.2}$$

We find the points of local extremum: $z=0$ and the roots of a quadratic equation

$$\Lambda\ \mathrm{ch}^4 z-2b\gamma\ \mathrm{ch}^2 z+\gamma^2\Lambda=0\qquad\Longrightarrow$$
$$\left(\mathrm{ch}^2 z\right)|_{1,2}=\frac{b}{\Lambda}\gamma\pm\sqrt{(\frac{b^2}{\Lambda^2}-1)\gamma^2}\ . \tag{4.3}$$

When considering the bound states (for motion in the variable r) we have $\Lambda^2 > b^2$. This means that the square root in (4.3) is an imaginary number. Consequently, the point of zero force (equilibrium points) except $z=0$ cannot exist. The situation is illustrated in the Fig. 1.

After the change of variables $\mathrm{ch}^2 z = y$, the differential equation (4.1) reads

$$\left[\frac{d^2}{dy^2}+\left(\frac{3}{2}\frac{1}{y}+\frac{1}{2}\frac{1}{y-1}\right)\frac{d}{dy}+\frac{\epsilon}{4y(y-1)}+\frac{b\gamma-\Lambda\ y}{(y-\gamma)(y+\gamma)4y(y-1)}\right]Z(y)=0\ . \tag{4.4}$$

Note that singular points $y=0,\pm\gamma(|\ \gamma\ |<<1)$ are located outside the physical range of the variable. Further progress in analytical treatment of eq. (4.4) (with 5 singular points) is hardly possible.



Figure 1: *Effective potential $U(z)$:*

5 Solution of the radial equation

Let us turn to the radial equation (3.7) for the function $R(r)$. It is solvable in hypergeometric functions – see more detail [4]. Below we will write done only final results on energy spectrum. There exist only finite series of bound states, defined by relations

$$m < 2b\ ,\ \ \frac{m+\mid m\mid}{2} + n + 1/2 \leq b\ ,$$
$$\Lambda - 1/4 = 2b\left(\frac{m+\mid m\mid}{2} + n + 1/2\right) - \left(\frac{m+\mid m\mid}{2} + n + \frac{1}{2}\right)^2\ ; \tag{5.1}$$

Λ obeys the restriction $b \leq \Lambda$. In usual units the last relation can be written as:

$$\Lambda - \frac{1}{4} = \rho^2\Lambda_0 - \frac{1}{4}\ ,\quad m < 2B,\quad m + n + 1/2 \leq \frac{eB}{\hbar c}\rho^2\ ,$$
$$\rho^2\Lambda_0 - \frac{1}{4} = 2\frac{eB}{\hbar c}\rho^2(\frac{m+\mid m\mid}{2} + n + 1/2) - (\frac{m+\mid m\mid}{2} + n + 1/2)^2, n = 0, 1, \ldots, N_B.$$

In the limit of vanishing curvature, we obtain the known result in the flat space

$$E - \frac{P^2}{2M} = \frac{eB\hbar}{Mc}\,(\frac{m+\mid m\mid}{2} + n + 1/2)\ .$$

6 Cox's particle in the electric field, Minkowski space

Schrödinger equation for Cox's particle in the electric field has the form (see Section **13**)

$$\left(D_t - c\,\frac{\Gamma^2 E_i E^i \mu + \Gamma E^j D_j}{2(1+\Gamma^2 E_i E^i)}\right)\Psi = -\frac{1}{2M}\,\overset{\circ}{D}_k\, g^{kj}\left[D_j + \frac{\Gamma^2 E_j (E^i D_i) + \mu\Gamma E_j}{1+\Gamma^2 E_i E^i}\right]\Psi\ ; \tag{6.1}$$

the notation is used:

$$A_0 = -eEz,\quad E_i = (F_{01}, F_{02}, F_{03})\ ,\quad g^{11}E_1 = E^1,\ g^{22}E_2 = E^2,\ g^{33}E_3 = E^3,$$
$$i\hbar\partial_t - eA_0 = D_t,\quad i\hbar\partial_k = D_k,\quad \frac{i\hbar}{\sqrt{-g}}\frac{\partial}{\partial x^k}\sqrt{-g} = \overset{\circ}{D}_k,\quad \mu = Mc\ .$$

Let us use cylindric coordinates

$$dS^2 = c^2dt^2 - dr^2 - r^2d\phi^2 - dz^2,\quad E_3 = E,\ E^3 = -E,\ E_3E^3 = -E^2. \tag{6.2}$$

First, we get (let it be $\Gamma E = \gamma$)

$$\left(D_t - c\frac{\Gamma^2 E_i E^i \mu + \Gamma E^j D_j}{2(1+\Gamma^2 E_i E^i)}\right) = i\hbar\partial_t + eEz + c\,\frac{\gamma^2\mu + \gamma D_3}{2(1-\gamma^2)}. \tag{6.3}$$

Next, we consider the Hamiltonian

$$H = \frac{1}{2M}\left[\overset{\circ}{D}_1\, D_1 + \overset{\circ}{D}_2\,\frac{1}{r^2}D_2 + \overset{\circ}{D}_3\left(\,D_3 + \frac{\mu\gamma}{1-\gamma^2}\,\right)\right]. \tag{6.4}$$

In explicit form, the extended Schrödinger equation looks as follows (to allow for the imaginary character of γ, we make formal change $i\gamma \longrightarrow \gamma$)

$$\left(i\hbar\partial_t + eEz - \frac{Mc^2\gamma^2}{2(1+\gamma^2)} + \frac{\gamma}{2(1+\gamma^2)}\hbar c\partial_z\right)\Psi$$
$$= \frac{-\hbar^2}{2M}\left(\partial_r^2 + \frac{1}{r}\partial_r + \frac{\partial_\phi^2}{r^2} + \partial_z^2 - \frac{(Mc/\hbar)\gamma}{1+\gamma^2}\partial_z\right)\Psi.$$

With the substitution $\Psi = e^{-iWt/\hbar}e^{im\phi}Z(z)R(t)$ and the notation

$$\frac{M^2c^2}{\hbar^2} = \frac{1}{\lambda^2}\ ,\ \frac{2M}{\hbar^2}W = w\ ,\ \frac{2M}{\hbar^2}\,eEz = \nu\ ,$$

we get

$$\frac{1}{Z(z)}\left(\partial_z^2 + \nu\ z + w - \frac{1}{\lambda^2}\frac{\gamma^2}{1+\gamma^2}\right)Z(z) + \frac{1}{R(r)}\left(\partial_r^2 + \frac{1}{r}\partial_r - \frac{m^2}{r^2}\right)R(r). \tag{6.5}$$

After separation of the variables ($w_\perp > 0$ stands for the separation constant) we derive

$$\left(\partial_r^2 + \frac{1}{r}\partial_r - \frac{m^2}{r^2} + w_\perp\right)R(r) = 0\ ,$$
$$(\partial_z^2 + \nu\ z + w')Z(z) = 0\ , \tag{6.6}$$

where

$$w' = w - w_\perp + \frac{1}{\lambda^2}\frac{\gamma^2}{1+\gamma^2} \ .$$

In fact, (6.6) coincide with the well known equations for an ordinary particle in the uniform electric field. Equation in the variable z looks as a one-dimensional Schrödinger equation in the potential of the form $U(z) = -\nu\ z\ ,\ \nu > 0$:

$$(\frac{d^2}{dz^2} + w' + \nu\ z)Z(z) = 0 \ . \tag{6.7}$$

The form of the curve $U(z)$ says that any particle moving from the right must be reflected by this barrier in vicinity of the point $z_0 = -\frac{w'}{\nu}$ (we assume that electric force acts in positive direction of the axis z).

Mathematic solutions of the equation (6.7) can be expressed in Airy function. Indeed, let us change the variable

$$\nu z + w' = ax,\ (\frac{d^2}{dx^2} + \frac{a^3}{\nu^2}x)Z(x) = 0 \ ;$$

let it be (for definiteness $\nu > 0$)

$$\frac{a^3}{\nu^2} = -1, \qquad a = -\nu^{2/3}, \quad x = \frac{\nu z + w'}{-\nu^{2/3}} = -\nu^{1/3}z - \frac{w'}{\nu^{2/3}}; \tag{6.8}$$

then we arrive at the Airy equation

$$\left(\frac{d^2}{dx^2} - x\right) Z(x) = 0 \ ; \tag{6.9}$$

to the turning point $z_0 = -w'/\nu$ there corresponds the value $x_0 = 0$.

Eq. (6.9) can be related to the Bessel equation. Indeed, let us introduce the variable

$$\xi = \frac{2}{3}x^{3/2} \ , \qquad x = \frac{3}{2}\xi^{2/3} \ ,$$

then Airy equation gives

$$(\frac{1}{3\xi}\frac{d}{d\xi} + \frac{d^2}{d\xi^2} - 1)Z = 0 \ .$$

Applying the substitution $Z = \xi^{1/3}f(\xi)$, we arrive at the Bessel equation [8]

$$\left(\frac{d^2}{d\xi^2} + \frac{1}{\xi}\frac{d}{d\xi} - 1 - \frac{1/9}{\xi^2}\right) f(\xi) = 0 \tag{6.10}$$

with two linearly independent solutions

$$f_1(\xi) = J_{+1/3}(i\xi) \ ,\ f_2(\xi) = J_{-1/3}(i\xi) \ . \tag{6.11}$$

Thus, general solutions of Airy equation can be constructed as linear combinations of

$$Z_1(x) = \xi^{1/3}J_{+1/3}(i\xi) \ ,\ \ Z_2(x) = \xi^{1/3}J_{-1/3}(i\xi) \ ,$$

where

$$i\xi = i\frac{2\sqrt{\nu}}{3}\ (z + \frac{w'}{\nu})^{3/2}\ .$$

With the use of the known relation [8, 8]

$$J_\mu(y) = \frac{(y/2)^\mu}{\Gamma(\mu+1)}\ e^{-iy}\ {}_1F_1(\mu+\frac{1}{2}, 2\mu+1, 2iy)$$

and with the notation $y = i\xi,\ \mu = +1/3, -1/3$, one expresses two independent solutions of the Schrödinger equation as follows

$$Z_1 = \xi^{1/3} J_{+1/3}(i\xi) = \xi^{+1/3}\frac{(i\xi/2)^\mu}{\Gamma(\mu+1)} e_1^\xi F_1(+\mu+\frac{1}{2}, +2\mu+1, -2\xi)\ ,$$
$$Z_2 = \xi^{1/3} J_{-1/3}(i\xi) = \xi^{1/3}\frac{(i\xi/2)^{-\mu}}{\Gamma(-\mu+1)}\ e_1^\xi F_1(-\mu+\frac{1}{2}, -2\mu+1, -2\xi)\ .$$

7 Cox's Particle in the electric field in the Lobachevsky model

We determine the concept of generalized electric field in the special system of cylindric coordinates in the curved space as follows [6]

$$dS^2 = dt^2 - \mathrm{ch}^2 z(dr^2 + \mathrm{sh}^2 r d\phi^2) - dz^2,$$
$$\sqrt{-g} = \mathrm{sh}\ r\ \mathrm{ch}^2 z,\ A_0 = -E\rho \tanh z,$$
$$E_3 = \frac{E}{\cosh^2 z},\ E^3 = -\frac{E}{\cosh^2 z},\ E_3 E^3 = -E^2\ \cosh^{-4} z. \tag{7.1}$$

Below we use operators (the coordinate $tc/\rho \longrightarrow t$ is dimensionless)

$$i\frac{\hbar c}{\rho}\partial_t - eA_0 = D_t,\ i\frac{\hbar}{\rho}\partial_k = D_k, \quad \frac{i\hbar/\rho}{\sqrt{-g}}\frac{\partial}{\partial x^k}\sqrt{-g} = \overset{\circ}{D}_k\ .$$

We start with the extended Schrödinger equation (let it be $E^2\ \cosh^{-4} z = -\gamma^2(x)$)

$$\left(D_t - c\ \frac{\Gamma^2 E_i E^i \mu + \Gamma E^j D_j}{2(1+\Gamma^2 E_i E^i)}\right)\ \Psi = \frac{1}{2M\rho^2}$$
$$= \overset{\circ}{D}_k\ (-g^{kj})\left(D_j + \frac{\Gamma^2 E_j(E^i D_i) + \mu\Gamma E_j}{1+\Gamma^2 E_i E^i}\right)\Psi.$$

Allowing for relations

$$\left(D_t - c\ \frac{\Gamma^2 E_i E^i \mu + \Gamma E^j D_j}{2(1+\Gamma^2 E_i E^i)}\right)$$
$$= \frac{\hbar c}{\rho}\left(i\partial_t + \frac{eE\rho}{\hbar c/\rho}\tanh z + \frac{1}{2}\frac{Mc\rho}{\hbar}\frac{\gamma^2(z)}{1-\gamma^2(z)} + \frac{1}{2}\frac{\gamma(z)}{1-\gamma^2(z)} i\partial_z\right),$$

and

$$H = -\frac{\hbar^2}{2M\rho^2}\left[\frac{1}{\cosh^2 z}\left(\partial_r^2 + \frac{\cosh r}{\sinh r}\partial_r + \frac{\partial_\phi^2}{\sinh^2 r}\right) + (\partial_z + 2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z})\left(\partial_z - \frac{Mc\rho}{\hbar}\frac{i\gamma(z)}{1-\gamma^2(z)}\right)\right],$$

we get e an explicit form of the extended Schrödinger equation

$$\frac{\hbar c}{\rho}\left(i\partial_t+\frac{eE\rho}{\hbar c/\rho}\tanh z+\frac{1}{2}\frac{Mc\rho}{\hbar}\frac{\gamma^2(z)}{1-\gamma^2(z)}+\frac{1}{2}\frac{\gamma(z)}{1-\gamma^2(z)}i\partial_z\right)\Psi$$
$$=-\frac{\hbar^2}{2M\rho^2}\left[\frac{1}{\cosh^2 z}\left(\partial_r^2+\frac{\cosh r}{\sinh r}\partial_r+\frac{\partial_\phi^2}{\sinh^2 r}\right)+(\partial_z+2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z})\left(\partial_z-\frac{Mc\rho}{\hbar}\frac{i\gamma(z)}{1-\gamma^2(z)}\right)\right]\Psi;$$

Specially note that two terms proportional to $i\gamma(z)\partial_z$ compensate each other. Additionally, we should perform formal change $i\gamma\longrightarrow\gamma$:

$$\frac{\hbar c}{\rho}\left(i\partial_t+\frac{eE\rho}{\hbar c/\rho}\tanh z-\frac{1}{2}\frac{Mc\rho}{\hbar}\frac{\gamma^2(z)}{1+\gamma^2(z)}\right)\Psi$$
$$=-\frac{\hbar^2}{2M\rho^2}\left[\frac{1}{\cosh^2 z}\left(\partial_r^2+\frac{\cosh r}{\sinh r}\partial_r+\frac{\partial_\phi^2}{\sinh^2 r}\right)\right.$$
$$\left.+(\partial_z+2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z})\partial_z-\frac{Mc\rho}{\hbar}\left(\frac{\partial}{\partial z}\frac{\gamma(z)}{1+\gamma^2(z)}\right)-\frac{Mc\rho}{\hbar}\frac{\gamma(z)}{1+\gamma^2(z)}2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z}\right]\Psi;$$

With the use of substitution

$$\Psi=\exp(-iwt)\ e^{im\phi}R(r)Z(z),\ \ w=\frac{W\rho}{\hbar c},$$

and notation

$$W=w\frac{\hbar c}{\rho}\frac{1}{\hbar^2/2M\rho^2}=2w\frac{M\rho c}{\hbar},\quad \nu=\frac{eE\rho}{\hbar^2/2M\rho^2}\ ,$$
$$\frac{1}{2}Mc^2\frac{1}{\hbar^2/2M\rho^2}=\frac{M^2\rho^2c^2}{\hbar^2}=\mu^2,$$

we get

$$\cosh^2 z\left(W+\nu\ \tanh z-\frac{\mu^2\gamma^2(z)}{1+\gamma^2(z)}\right)RZ+\left(\partial_r^2+\frac{\cosh r}{\sinh r}\partial_r-\frac{m^2}{\sinh^2 r}\right)RZ$$
$$+\cosh^2 z\left[(\partial_z+2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z})\partial_z-\mu\left(\frac{\partial}{\partial z}\frac{\gamma(z)}{1+\gamma^2(z)}\right)-\mu\frac{\gamma(z)}{1+\gamma^2(z)}2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z}\right]RZ=0\ . \tag{7.2}$$

In this equation one can separate the variables

$$\left(\frac{d^2}{dr^2}+\frac{\cosh r}{\sinh r}\frac{d}{dr}-\frac{m^2}{\sinh^2 r}+\Lambda\right)R=0, \tag{7.3}$$

$$\left[\frac{d^2}{dz^2}+2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z}\frac{d}{dz}-\mu\left(\frac{d}{dz}\frac{\gamma(z)}{1+\gamma^2(z)}\right)\right.$$
$$\left.-\mu\frac{\gamma(z)}{1+\gamma^2(z)}2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z}+W+\nu\ \tanh z-\mu^2\frac{\gamma^2(z)}{1+\gamma^2(z)}-\frac{\Lambda}{\mathrm{ch}^2 z}\right]Z=0\ ; \tag{7.4}$$

remember that $\gamma(z) = \gamma\ \mathrm{ch}^{-2} z$. The most interesting is the equation in variable z. After elementary transformation it is reduced to the form

$$\left(\frac{d^2}{dz^2} + 2\frac{\mathrm{sh}\ z}{\mathrm{ch}\ z}\frac{d}{dz}\right.$$
$$-2\mu\gamma\ \mathrm{sh}\ z\ \mathrm{ch}\ z\frac{-\mathrm{ch}^4 z + \gamma^2}{(\mathrm{ch}^4 z + \gamma^2)^2} - 2\mu\gamma\ \frac{\mathrm{sh}\ z\ \mathrm{ch}\ z}{\mathrm{ch}^4 z + \gamma^2}$$
$$\left. +W + \nu\ \tanh z - \frac{\mu^2\gamma^2}{\mathrm{ch}^4 z + \gamma^2} - \frac{\Lambda}{\mathrm{ch}^2 z}\right) Z) = 0\ . \tag{7.5}$$

This final equation turns out to be very complex and it hardly can be solved analytically.

8 Solving the radial equation in the Lobachewsky space

In eq. (7.3) let us introduce a new variable $x = (1 + \cosh r)/2,\ x \in [1,\ +\infty)$, so that

$$x\,(1-x)\,\frac{d^2R}{dx^2} + (1-2\,x)\,\frac{dR}{dx} - \left(w_\perp + \frac{1}{4}\,\frac{m^2}{x} + \frac{1}{4}\,\frac{m^2}{1-x}\right)\,R = 0\,; \tag{8.1}$$

with the substitution $R = x^a\,(1-x)^b\,F$ at

$$a = \pm\frac{|\,m\,|}{2}\,,\quad b = \pm\frac{|\,m\,|}{2}$$

we obtain the hypergeometric equation

$$x\,(1-x)\,\frac{d^2F}{dx^2} + [2\,a + 1 - (2\,a + 2\,b + 2)\,x]\,\frac{dF}{dx}\, - [(a+b)\,(a+b+1) + w_\perp]\,F = 0\,,$$

with parameters

$$F = F(\alpha, \beta, \gamma; x)\ ,$$
$$\alpha = a + b + \frac{1}{2} - i\sqrt{w_\perp - 1/4}\,,$$
$$\beta = a + b + \frac{1}{2} + i\sqrt{w_\perp - 1/4}\,,$$
$$w_\perp > \frac{1}{4}\,,\qquad \gamma = 2\,a + 1\,. \tag{8.2}$$

We will specify solutions tending to zero at $r = 0$:

$$F = u_2 = F(\alpha, \beta, \alpha + \beta + 1 - \gamma, 1 - x)\ ; \tag{8.3}$$

when a and b take positive values

$$a = +\frac{|\,m\,|}{2}\,,\qquad b = +\frac{|\,m\,|}{2}\,; \tag{8.4}$$

the complete radial function is

$$R = x^a(1-x)^b \qquad \times F(\alpha, \beta, \alpha + \beta + 1 - \gamma, 1 - x). \tag{8.5}$$

To find behavior at infinity, $r \to +\infty$, one should apply the following Kummer relationship [7]

$$u_2 = \frac{\Gamma(\alpha+\beta+1-\gamma)\Gamma(\beta-\alpha)}{\Gamma(\beta+1-\gamma)\Gamma(\beta)} e^{-i\pi\alpha}\ u_3 + \frac{\Gamma(\alpha+\beta+1-\gamma)\Gamma(\alpha-\beta)}{\Gamma(\alpha+1-\gamma)\Gamma(\alpha)} e^{-i\pi\beta}\ u_4\,, \tag{8.6}$$

where

$$u_2 = F(\alpha,\beta,\alpha+\beta+1-\gamma;1-x)\ ,$$
$$u_3 = (-x)^{-\alpha} F(\alpha,\alpha+1-\gamma,\alpha+1-\beta,\frac{1}{x})\ ,$$
$$u_4 = (-x)^{-\beta} F(\beta,\beta+1-\gamma,\beta+1-\alpha,\frac{1}{x})\ .$$

Therefore, asymptotic behavior at $x \to 1$ ($r \to +\infty$) is given by

$$R \approx (-1)^{a+b}\Gamma(\alpha+\beta+1-\gamma) \times \left(\frac{\Gamma(\beta-\alpha)}{\Gamma(\beta+1-\gamma)\Gamma(\beta)} e^{-i\pi\alpha}\ (-x)^{a+b-\alpha} + \frac{\Gamma(\alpha-\beta)}{\Gamma(\alpha+1-\gamma)\Gamma(\alpha)} e^{-i\pi\beta}\ (-x)^{a+b-\beta} \right). \tag{8.7}$$

From this it follows $x \approx \frac{e^r}{4}$ and

$$R \approx (-1)^{a+b}\Gamma(\alpha+\beta+1-\gamma)(-x)^{-1/2} \times \left(\frac{\Gamma(\beta-\alpha)}{\Gamma(\beta+1-\gamma)\Gamma(\beta)} e^{-i\pi\alpha}\ (-x)^{+i\sqrt{\lambda-1/4}} + \frac{\Gamma(\alpha-\beta)}{\Gamma(\alpha+1-\gamma)\Gamma(\alpha)} e^{-i\pi\beta}\ (-x)^{-i\sqrt{\lambda-1/4}} \right). \tag{8.8}$$

Thus, constructed solutions represent standing radial waves. The factor $e^{-r/2}$ is not significant for probability interpretation

$$dW = \sqrt{-g}\ \psi^*\psi\ ,$$

and the term $e^{-r/2}$ will be compensated by the factor $\sinh r \approx e^{+r}/2$ entering the volume element $dV = \sqrt{-g}\ dr\ dz\ d\phi$.

9 Cox's particle in magnetic field in the spherical Riemann space

In cylindric coordinates of the spherical Riemann space (below ρ stands for the curvature radius)

$$dS^2 = dt^2 - \cos^2 z\,(dr^2 + \sin^2 r\, d\phi^2) - dz^2, \quad \sqrt{-g} = \sin\ r\ \cos^2 z\,,$$

analogue of the uniform magnetic field is given by relations – see [4, 5]:

$$A_\phi = B\rho^2\,(\cos\ r - 1)\,,\ F_{r\phi} = B\rho\,\sin\ r\,,$$
$$B_3 = B\rho\sin r,\ B^3 = \frac{B}{\rho\sin r\cos^4 z}, B_iB^i = \frac{B^2}{\cos^4 z}.$$

We start with extended Schrödinger equation in the form

$$D_t\,\Psi = \frac{1}{2M\rho^2}\left[\overset{\circ}{D}_1\,\frac{1}{\cos^2 z}\,\overset{*}{D}_1 + \overset{\circ}{D}_2\,\frac{1}{\sin^2 r\,\cos^2 z}\,\overset{*}{D}_2 + \overset{\circ}{D}_3\overset{*}{D}_3\right]\Psi\,, \qquad (9.1)$$

where

$$D_1 = i\hbar\partial_r\,, \qquad D_2 = i\hbar\partial_\phi - \frac{e}{c}\,B\rho^2(\cos\,r-1)\,, \qquad D_3 = i\hbar\partial_z\,,$$

$$\overset{\circ}{D}_1 = i\hbar(\partial_r + \frac{\cos\,r}{\sin\,r})\,, \qquad \overset{\circ}{D}_2 = i\hbar\partial_\phi - \frac{e}{c}B\rho^2(\cos\,r-1)\,, \qquad \overset{\circ}{D}_3 = i\hbar(\partial_z - 2\frac{\sin\,z}{\cos\,z})\,,$$

$$\overset{*}{D}_1 = \frac{(D_1 - \Gamma B_3 D^2)}{1+\Gamma^2B^2\cos^{-4}z} = \frac{1}{1+\Gamma^2B^2\cos^{-4}z}\left[i\hbar\partial_r + \Gamma B\,\sin r\,\cos^2 z\left(i\hbar\partial_\phi - \frac{e}{c}B\rho^2(\cos\,r-1)\right)\right],$$

$$\overset{*}{D}_2 = \frac{(\,D_2 + \Gamma B_3 D^1)}{1+\Gamma^2B^2\cos^{-4}z} = \frac{1}{1+\Gamma^2B^2\cos^{-4}z}\left[i\hbar\partial_\phi - \frac{e}{c}B\rho^2(\cos\,r-1) - i\hbar\,\frac{\Gamma B\sin r}{\cos^2 z}\,\partial_r\right],$$

$$\overset{*}{D}_3 = \frac{1}{1+\Gamma^2B^2\cos^{-4}z}(D_3 + \Gamma^2B^3\,B_3D_3) = i\hbar\partial_z\,.$$

With notation $\frac{eB\rho^2}{\hbar c} = b$, $\frac{\Gamma B}{\cos^2 z} = \gamma(z)$, we get

$$\frac{1}{2M\rho^2}\,\overset{\circ}{D}_1\,g^{11}\,\overset{*}{D}_1$$

$$= \frac{\hbar^2}{2M\rho^2\cos^2 z\,(1+\gamma^2(z))}\left[\partial_r^2 + \frac{\cos r}{\sin r}\,\partial_r + i\gamma(z)b\frac{\cos r - 1}{\sin r}\,\partial_r + \frac{\gamma(z)}{\sin r}\partial_r\partial_\phi - i\gamma(z)b\right],$$

$$\frac{1}{2M\rho^2}\,\overset{\circ}{D}_2\,g^{22}\,\overset{*}{D}_2$$

$$= \frac{\hbar^2}{2M\rho^2\sin^2 r\,\cos^2 z\,(1+\gamma^2(z))}\,[\partial_\phi + ib(\cos\,r-1)]\,[\partial_\phi + ib(\cos\,r-1) - \gamma(z)\sin r\,\partial_r]\,,$$

$$\frac{1}{2M\rho^2}\,\overset{\circ}{D}_3\,g^{33}\,\overset{*}{D}_3 = \frac{\hbar^2}{2M\rho^2}\,(\partial_z - 2\frac{\sin\,z}{\cos\,z})\,\partial_z\,.$$

Using the substitution $\Psi = e^{-iEt/\hbar}e^{im\phi}Z(z)R(r)$, $\epsilon = E/(\hbar^2/2M\rho^2)$, we reduce the Schrödinger equation to the form (the function $\gamma(z)$ must be imaginary: $i\gamma(z) \Longrightarrow \gamma(z)$)

$$\left[\frac{1}{\cos^2 z\,(1-\gamma^2(z))}\left(\partial_r^2 + \frac{\cos\,r}{\sin\,r}\,\partial_r - \frac{[m+b(\cos\,r-1)]^2}{\sin^2 r} - b\gamma(z)\right)\right.$$

$$\left. +\epsilon + (\partial_z - 2\frac{\sin\,z}{\cos\,z})\partial_z\right]R(r)Z(z) = 0\,.$$

One separates the variables

$$\left[\frac{d^2}{dr^2} + \frac{\cos\,r}{\sin\,r}\frac{d}{dr} - \frac{[m+b(\cos\,r-1)]^2}{\sin^2 r} + \Lambda\right]R(r) = 0\,, \qquad (9.2)$$

$$\left(\frac{d^2}{dz^2} - 2\,\frac{\sin\,z}{\cos\,z}\frac{d}{dz} + \epsilon - \frac{b\gamma + \Lambda\,\cos^2 z}{\cos^4 z - \gamma^2}\right)Z(z) = 0\,. \qquad (9.3)$$

10 Analysis of the equation in the variable z

Excluding in (9.3) the term with first derivative, we obtain an equation in the effective potential

$$Z(z) = \frac{1}{\cos\, z} f(z)\,, \quad \left(\frac{d^2}{dz^2} + \epsilon + 1 - U(z)\right) f(z) = 0\,,$$
$$U(z) = \frac{b\gamma + \Lambda\, \cos^2 z}{\cos^4 z - \gamma^2}\,, \quad U(z=0) = \frac{b\gamma + \Lambda}{1-\gamma^2}\,, \quad U(z = \pm\frac{\pi}{2}) = -\frac{b}{\gamma}\,. \tag{10.1}$$

Expression for an effective force looks

$$F_z = -\frac{dU}{dz} = -2\, \cos\, z\, \sin\, z\, \frac{\Lambda\, \cos^4 z + 2\, b\, \gamma\, \cos^2 z + \gamma^2 \Lambda}{(\cos^4 z - \gamma^2)^2}\,; \tag{10.2}$$

points of vanishing force (or of local extremum) are $z = 0$ and the roots of the quadratic equation

$$\Lambda\, \cos^4 z + 2\, b\, \gamma\, \cos^2 z + \gamma^2 \Lambda = 0 \qquad \Longrightarrow \qquad \left(\cos^2 z\right)|_{1,2} = -\frac{b}{\Lambda}\gamma \pm \sqrt{(\frac{b^2}{\Lambda^2} - 1)\gamma^2}\,. \tag{10.3}$$

Due to inequality $\Lambda^2 > b^2$, under the square root is imaginary quantity. Therefore, in physical region of the variable z we have no other force vanishing point in addition to $z = 0$. In the new variable $\cos^2 z = y$, differential equation (9.3) takes the form

$$\left[\frac{d^2}{dy^2} + \left(\frac{3}{2}\frac{1}{y} + \frac{1}{2}\frac{1}{y-1}\right)\frac{d}{dy} - \frac{\epsilon}{4\, y\, (y-1)} + \frac{b\gamma + \Lambda\, y}{(y-\gamma)\,(y+\gamma)\, 4\, y\, (y-1)}\right] Z(y) = 0\,. \tag{10.4}$$

Behavior near the five singular pints can be found straightforwardly.

$y \sim 1 \;\; (z \to 0)$

$$\left[\frac{d^2}{dy^2} + \left(\frac{1}{2}\frac{1}{y-1}\right)\frac{d}{y} - \frac{\epsilon}{4(y-1)} + \frac{b\gamma + \Lambda}{(1-\gamma^2)\, 4\, (y-1)}\right] Z(y) = 0\,,$$
$$Z(y) = \exp\left[\pm\sqrt{A(y-1)}\right], \qquad A = \epsilon - \frac{b\gamma + \Lambda}{1-\gamma^2}\,; \tag{10.5}$$

$y \sim 0$

$$\left(\frac{d^2}{dy^2} + \frac{3}{2y}\,\frac{d}{dy} + \frac{\epsilon}{4y} + \frac{b}{4\gamma y}\right) Z(y) = 0\,,$$
$$Z(y) = \frac{\exp[\,\pm\sqrt{Cy}\,]}{\sqrt{y}}, \quad C = -\epsilon - \frac{b}{\gamma}\,; \tag{10.6}$$

$y \sim \infty$

$$\left(\frac{d^2}{dy^2} + \frac{2}{y}\frac{d}{dy} - \frac{\epsilon}{4y^2}\right) Z(z) = 0, \quad Z = y^D\,, \quad D = \frac{-1 \pm \sqrt{\epsilon + 1}}{2}\,; \tag{10.7}$$

$y \sim +\gamma$

$$\left[\frac{d^2}{dy^2} + \frac{1}{2}\left(\frac{3}{\gamma} + \frac{1}{\gamma - 1}\right)\frac{d}{dy} + \frac{\Lambda + b}{8\gamma(1-\gamma)}\;\; \frac{1}{y-\gamma}\right] Z(y) = 0$$

or

$$\left[(y-\gamma)\frac{d^2}{dy^2}+M(y-\gamma)\frac{d}{dy}+N\right]Z(y)=0,\quad M=\frac{1}{2}\left(\frac{3}{\gamma}+\frac{1}{\gamma-1}\right),\quad N=\frac{\Lambda+b}{8\gamma(1-\gamma)}\,.$$

Changing the variable $-M(y-\gamma)=x$ we get

$$\left(x\frac{d^2}{dx^2}-x\frac{d}{dx}-\alpha\right)Z=0,\qquad \alpha=\frac{N}{M}\,,$$

which is confluent hypergeometric equation of a special form

$$\left(x\frac{d^2}{dx^2}+(c-x)\frac{d}{dx}-a\right)Z=0\,,\qquad c=0,\quad a=\frac{N}{M}=\frac{\Lambda+b}{4(3-4\gamma)}\,;$$

its general solution looks as $Z=c_1M(a+1,2,y)+c_2U(a+1,2,y)$.

Now, let us consider the case $y\sim-\gamma$:

$$\left[\frac{d^2}{dy^2}\frac{1}{2}\left(\frac{3}{-\gamma}+\frac{1}{-\gamma-1}\right)\frac{d}{dy}+\frac{\Lambda-b}{8(-\gamma)(1+\gamma)}\,\frac{1}{y+\gamma}\right]Z(y)=0$$

or shorter

$$\left[(y-\gamma)\frac{d^2}{dy^2}+M'(y-\gamma)\frac{d}{dy}+N'\right]Z(y)=0,\ -M'(y+\gamma)=x\,,$$

$$\left(x\frac{d^2}{dx^2}-x\frac{d}{dx}-\alpha'\right)Z=0,\ a'=\frac{N'}{M'}\,,$$

which is confluent hypergeometric equation of a special form

$$\left(x\frac{d^2}{dx^2}+(c-x)\frac{d}{dx}-a\right)Z=0=0\,,\ c=0,\quad \alpha=\frac{N'}{M'}=\frac{\Lambda-b}{4(3+4\gamma)}\,;$$

its general solution is $Z=c_1M(a+1,2,y)+c_2U(a+1,2,y)$.

Further analytical treatment of the differential equation (10.4) is very difficult because complexity of this equation.

11 Solutions of the radial equation

Now let us consider the radial equation (9.2). It can be solved in hypergeometric functions – see [4, 5]. Below we will write down only energy spectrum. In usual measure units these formulas read

$$\begin{aligned}
m>0,\quad \rho^2\Lambda_0+\frac{1}{4}&=+2\frac{eB}{\hbar c}\rho^2(n+m+1/2)+(n+m+1/2)^2\,;\\
m<-2\frac{eB}{\hbar c}\,\rho^2,\quad \rho^2\Lambda_0+\frac{1}{4}&=-2\frac{eB}{\hbar c}\rho^2(n-m+1/2)+(n-m+1/2)^2\,;\\
-2\frac{eB}{\hbar c}\,\rho^2<m\le 0\,,\quad \rho^2\Lambda_0+\frac{1}{4}&=2\frac{eB}{\hbar c}\rho^2(n+1/2)+(n+1/2)^2\,.
\end{aligned}\qquad(11.1)$$

Transition to the limit of the Minkowski space is achieved accordingly to ($\rho\to\infty$)

$$m<0\,,\qquad \Lambda_0=2\frac{eB}{\hbar c}(n+1/2)\,;\qquad m\ge 0\,,\qquad \Lambda_0=+2\frac{eB}{\hbar c}(n+m+1/2)\,,\qquad(11.2)$$

Thus, we get the well-known result

$$E-\frac{P^2}{2M}=\ \frac{eB\hbar}{Mc}\,(\frac{m+\mid m\mid}{2}+n+1/2)\,.\qquad(11.3)$$

12 Cox's particle in electric field in the spherical model

In cylindrical coordinate $dS^2 = dt^2 - \cos^2 z\,(dr^2 + \sin^2 r\, d\phi^2) - dz^2$, external electric field along the axis z is given by

$$A_0 = -E\,\rho\,\tan z\,, \quad E_3 = \frac{E}{\cos^2 z}\,, \quad E^3 = -\frac{E}{\cos^2 z}\,, \;\; E_3\,E^3 = -\frac{E^2}{\cos^4 z}\,. \tag{12.1}$$

Below we use operators (and dimensionless coordinate $tc/\rho \longrightarrow t$)

$$i\frac{\hbar c}{\rho}\partial_t - eA_0 = D_t,\; i\frac{\hbar}{\rho}\partial_k = D_k, \frac{i\hbar/\rho}{\sqrt{-g}}\frac{\partial}{\partial x^k}\sqrt{-g} = \overset{\circ}{D}_k\,.$$

We start with the form of Schrödinger equation

$$\left(D_t - c\,\frac{\Gamma^2 E_i E^i \mu + \Gamma E^j D_j}{2(1+\Gamma^2 E_i E^i)}\right)\Psi = \frac{\overset{\circ}{D}_k\,(-g^{kj})}{2M\rho^2}\left(D_j + \frac{\Gamma^2 E_j (E^i D_i) + \mu\Gamma E_j}{1+\Gamma^2 E_i E^i}\right)\Psi. \tag{12.2}$$

After needed calculation we get representation for the wave equation

$$\frac{\hbar c}{\rho}\left(i\partial_t + \frac{eE\rho}{\hbar c/\rho}\tan z + \frac{1}{2}\frac{Mc\rho}{\hbar}\frac{\gamma^2(z)}{1-\gamma^2(z)} + \frac{1}{2}\frac{\gamma(z)}{1-\gamma^2(z)}i\partial_z\right)\Psi$$

$$= -\frac{\hbar^2}{2M\rho^2}\left[\frac{1}{\cos^2 z}\left(\partial_r^2 + \frac{\cos r}{\sin r}\partial_r + \frac{\partial_\phi^2}{\sin^2 r}\right) + \left(\partial_z - 2\,\frac{\sin z}{\cos z}\right)\left(\frac{1-2\gamma^2(z)}{1-\gamma^2(z)}\partial_z - \frac{Mc\rho}{\hbar}\frac{i\gamma(z)}{1-\gamma^2(z)}\right)\right]\Psi.$$

Note that two terms proportional to $i\gamma(z)\partial_z$ compensate each other; besides we make the formal change $i\gamma \longrightarrow \gamma$:

$$\frac{\hbar c}{\rho}\left(i\partial_t + \frac{eE\rho}{\hbar c/\rho}\tan z - \frac{1}{2}\frac{Mc\rho}{\hbar}\frac{\gamma^2(z)}{1+\gamma^2(z)}\right)\Psi$$

$$= -\frac{\hbar^2}{2M\rho^2}\left[\frac{1}{\cos^2 z}\left(\partial_r^2 + \frac{\cos r}{\sin r}\,\partial_r + \frac{\partial_\phi^2}{\sin^2 r}\right) + \left(\partial_z - 2\,\frac{\sin\,z}{\cos\,z}\right)\left(\frac{1+2\,\gamma^2(z)}{1+\gamma^2(z)}\,\partial_z\right)\right.$$

$$\left. - \frac{Mc\rho}{\hbar}\,\frac{\partial}{\partial z}\frac{\gamma(z)}{1+\gamma^2(z)} + \frac{Mc\rho}{\hbar}\frac{\gamma(z)}{1+\gamma^2(z)}\,2\,\frac{\sin\,z}{\cos\,z}\right]\Psi. \tag{12.3}$$

With substitution $\Psi = e^{-iwt}\,e^{im\phi}\,R(r)\,Z(z)\,,\; w = \frac{W\rho}{\hbar c}$, and notation

$$W = w\frac{\hbar c}{\rho}\,\frac{1}{\hbar^2/2M\rho^2} = 2w\frac{M\rho c}{\hbar}\,, \quad \nu = eE\rho\,\frac{1}{\hbar^2/2M\rho^2}\,, \quad \frac{1}{2}Mc^2\,\frac{1}{\hbar^2/2M\rho^2} = \frac{M^2\rho^2c^2}{\hbar^2}\;\; = \mu^2\,,$$

we arrive at

$$\cos^2 z\left(W + \nu\,\tan z - \mu^2\frac{\gamma^2(z)}{1+\gamma^2(z)}\right)R(r)Z(z)$$

$$+\left(\partial_r^2 + \frac{\cos r}{\sin r}\,\partial_r - \frac{m^2}{\sin^2 r}\right)R(r)Z(z) + \cos^2 z\left[\left(\partial_z - 2\,\frac{\sin\,z}{\cos\,z}\right)\left(\frac{1+2\,\gamma^2(z)}{1+\gamma^2(z)}\,\partial_z\right)\right.$$

$$\left. -\mu\,\frac{\partial}{\partial z}\,\frac{\gamma(z)}{1+\gamma^2(z)} + \mu\frac{\gamma(z)}{1+\gamma^2(z)}\,2\,\frac{\sin\,z}{\cos\,z}\right]R(r)Z(z) = 0\,. \tag{12.4}$$

After the separation of the variable we obtain

$$\left(\frac{d^2}{dr^2}+\frac{\cos r}{\sin r}\frac{d}{dr}-\frac{m^2}{\sin^2 r}+\Lambda\right)R(r)=0\ , \tag{6.8}$$

this equation can be readily solved in hypergeometric functions. In turn, equation in z variable is

$$\left[\left(\frac{d}{dz}-2\,\frac{\sin\,z}{\cos\,z}\right)\left(\frac{1+2\,\gamma^2(z)}{1+\gamma^2(z)}\,\frac{d}{dz}\right)-\mu\,\frac{d}{dz}\,\frac{\gamma(z)}{1+\gamma^2(z)}+\mu\frac{\gamma(z)}{1+\gamma^2(z)}\,2\,\frac{\sin\,z}{\cos\,z}\right.$$
$$\left.+W+\nu\,\tan z-\mu^2\frac{\gamma^2(z)}{1+\gamma^2(z)}-\frac{\Lambda}{\cos^2 z}\right]Z(z)=0\ . \tag{12.5}$$

Remember that $\gamma(z)=\gamma\,\cos^{-2}z$. The last equation can be translated to the following form

$$\left(\frac{\cos^4 z+2\,\gamma^2}{\cos^4 z+\gamma^2}\,\frac{d^2}{dz^2}-2\,\frac{\sin\,z}{\cos\,z}\,\frac{\gamma^2\,\cos^4 z+2\,\gamma^4+\cos^8 z}{(\cos^4 z+\gamma^2)^2}\,\frac{d}{dz}-\mu\,\gamma\,\frac{\cos^2 z}{\cos^4 z+\gamma^2}\,\frac{d}{dz}\right.$$
$$\left.+4\,\mu\,\gamma^3\,\frac{\sin\,z\,\cos\,z}{(\cos^4 z+\gamma^2)^2}+W+\nu\,\tan z-\frac{\mu^2\gamma^2}{\cos^4 z+\gamma^2}-\frac{\Lambda}{\cos^2 z}\right)Z(z)=0\ . \tag{12.6}$$

We could not proceed further with this differential equation because of its complexity.

13 Cox's particle in arbitrary curved space time, general analysis

First, we will use a Proca-like generalized system in Cartesian coordinates in Minkowski space (note the notation: $\mu=Mc$)

$$(\mu\delta^\beta_\alpha+\lambda F_\alpha{}^\beta)\Phi_\beta=D_\alpha\Phi\ ,\ D^\alpha\Phi_\alpha=\mu\Phi\ ,$$

or shorter

$$\Lambda_\alpha{}^\beta\Phi_\beta=D_\alpha\Phi\ ,\ D^\alpha\Phi_\alpha=\mu\Phi\ . \tag{13.1}$$

The first equation in (13.1) can be multiplied by the inverse matrix $(\Lambda^{-1})_\rho{}^\alpha$, so we get

$$\Phi_\rho=(\Lambda^{-1})_\rho{}^\alpha D_\alpha\Phi\ ,\ D^\rho\Phi_\rho=\mu\Phi\ . \tag{13.2}$$

From (13.2), one can derive a generalized Klein–Fock–Gordon equation for the scalar function Φ:

$$\left[\ \mu D^\rho(\Lambda^{-1})_\rho{}^\alpha D_\alpha\ -\ \mu^2\right]\Phi=0\ . \tag{13.3}$$

Equation (13.3) can be rewritten in the form

$$\{\mu(\Lambda^{-1})_\rho{}^\alpha\ D^\rho D_\alpha+\mu\,[\ i\hbar\partial^\rho(\Lambda^{-1})_\rho{}^\alpha]\ D_\alpha-\mu^2\}\Phi=0\ . \tag{13.4}$$

We are to find an explicit form of the inverse matrix Λ^{-1} (the calculation of the inverse matrix performed in this section is valid only in Cartesian coordinates of the flat space; generalization to the case of a curved space or curvilinear coordinates in the flat space will be given below):

$$\Lambda=(\Lambda_\alpha{}^\beta)=\left|\begin{array}{cccc}\mu & -e_1 & -e_2 & -e_3\\ -e_1 & \mu & -b_3 & b_2\\ -e_2 & b_3 & \mu & -b_1\\ -e_3 & -b_2 & b_1 & \mu\end{array}\right|,\quad e_i=\lambda E_i,\ b_i=\lambda B_i. \tag{13.5}$$

The inverse matrix is defined by the formula

$$\Lambda^{-1} = \frac{1}{\det \Lambda} \begin{vmatrix} M_0{}^0 & -M_1{}^0 & +M_2{}^0 & -M_3{}^0 \\ -M_0{}^1 & +M_1{}^1 & -M_2{}^1 & +M_3{}^1 \\ +M_0{}^2 & -M_1{}^2 & +M_2{}^2 & -M_3{}^2 \\ -M_0{}^3 & +M_1{}^3 & -M_2{}^3 & +M_3{}^3 \end{vmatrix} .$$

The determinant of the matrix Λ equals to $\det \Lambda = \mu^4 - \mu^2 \, (\vec{e}^{\,2} - \vec{b}^{\,2}) - (\vec{e}\,\vec{b})^2$, and the minors are

$$M_0{}^0 = \mu^3 + \mu \, \vec{b}^{\,2} \, , \quad M_1{}^1 = \mu^3 + \mu(b_1^2 - e_2^2 - e_3^2) \, ,$$

$$M_2{}^2 = \mu^3 + \mu(b_2^2 - e_1^2 - e_3^2) \, , \quad M_3{}^3 = \mu^3 + \mu(b_3^2 - e_1^2 - e_2^2) \, ,$$

$$M_0{}^1 = -\mu^2 e_1 - \mu(e_2 b_3 - e_3 b_2) - b_1(\vec{e}\,\vec{b}) \, , \quad M_1{}^0 = -\mu^2 e_1 + \mu(e_2 b_3 - e_3 b_2) - b_1(\vec{e}\,\vec{b}) \, ,$$

$$M_0{}^2 = \mu^2 e_2 + \mu(e_3 b_1 - e_1 b_3) + b_2(\vec{e}\,\vec{b}) \, , \quad M_2{}^0 = \mu^2 e_2 - \mu(e_3 b_1 - e_1 b_3) + b_2(\vec{e}\,\vec{b}) \, ,$$

$$M_0{}^3 = -\mu^2 e_3 - \mu(e_1 b_2 - e_2 b_1) - b_3(\vec{e}\,\vec{b}) \, , \quad M_3{}^0 = -\mu^2 e_3 + \mu(e_1 b_2 - e_2 b_1) - b_3(\vec{e}\,\vec{b}) \, ,$$

$$M_1{}^2 = \mu^2 b_3 - \mu(e_1 e_2 + b_1 b_2) - e_3(\vec{e}\,\vec{b}) \, , \quad M_2{}^1 = -\mu^2 b_3 - \mu(e_1 e_2 + b_1 b_2) + e_3(\vec{e}\,\vec{b}) \, ,$$

$$M_1{}^3 = \mu^2 b_2 + \mu(e_1 e_3 + b_1 b_3) - e_2(\vec{e}\,\vec{b}) \, , \quad M_3{}^1 = -\mu^2 b_2 + \mu(e_1 e_3 + b_1 b_3) + e_2(\vec{e}\,\vec{b}) \, ,$$

$$M_2{}^3 = \mu^2 b_1 - \mu(e_2 e_3 + b_2 b_3) - e_1(\vec{e}\,\vec{b}) \, , \quad M_3{}^2 = -\mu^2 b_1 - \mu(e_2 e_3 + b_2 b_3) + e_1(\vec{e}\,\vec{b}) \, .$$

Let us detail two simple cases. The first is the presence of the electric field:

$$B_i = 0 \, , \qquad \Lambda = \begin{vmatrix} \mu & -e_1 & -e_2 & -e_3 \\ -e_1 & \mu & 0 & 0 \\ -e_2 & 0 & \mu & 0 \\ -e_3 & -0 & 0 & \mu \end{vmatrix} ,$$

$$\Lambda^{-1} = \frac{1}{\mu^4 - \mu^2 \vec{e}^{\,2}} \begin{vmatrix} \mu^3 & \mu^2 e_1 & \mu^2 e_2 & \mu^2 e_3 \\ \mu^2 e_1 & \mu^3 - \mu(e_2^2 + e_3^2) & \mu e_1 e_2 & \mu e_1 e_3 \\ \mu^2 e_2 & \mu e_1 e_2 & \mu^3 - \mu(e_1^2 + e_3^2) & \mu e_2 e_3 \\ \mu^2 e_3 & \mu e_1 e_3 & \mu e_1 e_3 & \mu^3 - \mu(e_1^2 + e_3^2) \end{vmatrix} .$$

The second is the case of the magnetic field:

$$E_i = 0, \Lambda = \begin{vmatrix} \mu & 0 & 0 & 0 \\ 0 & \mu & -b_3 & b_2 \\ 0 & b_3 & \mu & -b_1 \\ 0 & -b_2 & b_1 & \mu \end{vmatrix} , \qquad \Lambda^{-1} = \frac{1}{\mu^4 + \mu^2 \vec{b}^{\,2}}$$

$$\times \begin{vmatrix} \mu^3 + \mu \vec{b}^{\,2} & 0 & 0 & 0 \\ 0 & \mu^3 + \mu b_1^2 & \mu^2 b_3 + \mu b_1 b_2 & -\mu^2 b_2 + \mu b_1 b_3 \\ 0 & -\mu^2 b_3 + \mu b_1 b_2 & \mu^3 + \mu b_2^2 & \mu^2 b_1 + \mu b_2 b_3 \\ 0 & \mu^2 b_2 + \mu b_1 b_3 & -\mu^2 b_1 + \mu b_2 b_3 & \mu^3 + \mu b_3^2 \end{vmatrix} .$$

Now, let us derive an explicit form of the generalized scalar equation in the case of the uniform magnetic field:

$$K^{\rho\alpha} = g^{\rho\beta}(\Lambda^{-1})_{\beta}{}^{\alpha} = \frac{1}{\mu^4 + \mu^2 \vec{b}^{\,2}}$$
$$\times \begin{vmatrix} \mu^3 + \mu\vec{b}^{\,2} & 0 & 0 & 0 \\ 0 & -\mu^3 - \mu b_1^2 & -\mu^2 b_3 - \mu b_1 b_2 & \mu^2 b_2 - \mu b_1 b_3 \\ 0 & \mu^2 b_3 - \mu b_1 b_2 & -\mu^3 - \mu b_2^2 & -\mu^2 b_1 - \mu b_2 b_3 \\ 0 & -\mu^2 b_2 - \mu b_1 b_3 & \mu^2 b_1 - \mu b_2 b_3 & -m^3 - \mu b_3^2 \end{vmatrix} . \tag{13.6}$$

We are to specify the equation (13.3)

$$(K^{\rho\alpha} D_\rho D_\alpha - \mu)\,\Phi = 0\,, \qquad (D_\alpha) = (i\hbar\partial_0,\ i\hbar\partial_j - \frac{e}{c}A_j) \tag{13.7}$$

in the field $\vec{A} = \frac{1}{2}\vec{x} \times \vec{B}$. After simple calculation we arrive at (for brevity we use the parameter $\lambda/\mu = \Gamma$)

$$\Big[\ (1 + \Gamma^2 \vec{B}^{\,2}) D_0^2 - \vec{D}^{\,2} + i\hbar\frac{e}{c}\Gamma\vec{B}^{\,2} - \Gamma^2(\vec{B}\vec{D})^2 - \mu^2(1 + \Gamma^2\vec{B}^{\,2})\ \Big]\,\Phi = 0\,. \tag{13.8}$$

Now let us consider the case of homogeneous electric field:

$$A_0 = -\vec{E}\,\vec{x},\ \bar{K}^{\rho\alpha} = \frac{1}{\mu^4 - \mu^2\vec{e}^{\,2}}$$
$$\times \begin{vmatrix} \mu^3 & \mu^2 e_1 & \mu^2 e_2 & \mu^2 e_3 \\ -\mu^2 e_1 & -\mu^3 + \mu(e_2^2 + e_3^2) & -\mu e_1 e_2 & -\mu e_1 e_3 \\ -\mu^2 e_2 & -\mu e_1 e_2 & -\mu^3 + \mu(e_1^2 + e_3^2) & -\mu e_2 e_3 \\ -\mu^2 e_3 & -\mu e_1 e_3 & -\mu e_2 e_3 & -\mu^3 + \mu(e_1^2 + e_2^2) \end{vmatrix} . \tag{13.9}$$

The equation (13.3) will take the form

$$\Big[\ D_0^2 - (1 - \Gamma^2\vec{E}^{\,2})\vec{D}^{\,2} - i\hbar\frac{e}{c}\Gamma\vec{E}^{\,2} - \Gamma^2(\vec{E}\vec{D})^2 - \mu^2(1 - \Gamma^2\vec{E}^{\,2})\ \Big]\,\Phi = 0\,. \tag{13.10}$$

Let us construct the inverse matrix Λ^{-1} in tensor notation. We start from the explicit form of the tensor $(F_\alpha{}^\beta)$:

$$(F_\alpha{}^\beta) = \begin{vmatrix} 0 & -E_1 & -E_2 & -E_3 \\ -E_1 & 0 & -B_3 & B_2 \\ -E_2 & B_3 & 0 & -B_1 \\ -E_3 & -B_2 & B_1 & 0 \end{vmatrix}, \tag{13.11}$$

then compute the convolution of two tensors (in this section we perform calculations, based on Cartesian coordinates in Minkowski space)

$$(F_\alpha{}^\beta)(F_\beta{}^\rho) = \begin{vmatrix} \vec{E}^{\,2} & -E_2B_3 + E_3B_2 & E_1B_3 - E_3B_1 & -E_1B_2 + E_2B_1 \\ E_2B_3 - E_3B_2 & E_1^2 - B_2^2 - B_3^2 & E_1E_2 + B_1B_2 & E_1E_3 + B_1B_3 \\ -E_1B_3 + E_3B_1 & E_1E_2 + B_1B_2 & E_2^2 - B_1^2 - B_3^2 & E_2E_3 + B_2B_3 \\ E_1B_2 - E_2B_1 & E_1E_3 + B_1B_3 & E_2E_3 + B_2B_3 & E_3^2 - B_1^2 - B_2^2 \end{vmatrix} . \tag{13.12}$$

Compute the convolution of three tensors:

$$(F_\alpha{}^\beta)(F_\beta{}^\rho)(F_\rho{}^\sigma) = (\vec{E}^{\,2} - \vec{B}^{\,2}) \begin{vmatrix} 0 & -E_1 & -E_2 & -E_3 \\ -E_1 & 0 & -B_3 & B_2 \\ -E_2 & B_3 & 0 & -B_1 \\ -E_3 & -B_2 & B_1 & 0 \end{vmatrix}$$
$$+(\vec{E}\vec{B}) \begin{vmatrix} 0 & -B_1 & -B_2 & -B_3 \\ -B_1 & 0 & E_3 & -E_2 \\ -B_2 & -E_3 & 0 & E_1 \\ -B_3 & E_2 & -E_1 & 0 \end{vmatrix} . \tag{13.13}$$

Next compute the convolution of the electromagnetic tensor and tensor dual to it:

$$(F_\alpha^{\times\beta})(F_\beta{}^\rho) = \begin{vmatrix} \vec{E}\vec{B} & 0 & 0 & 0 \\ 0 & \vec{E}\vec{B} & 0 & 0 \\ 0 & 0 & \vec{E}\vec{B} & 0 \\ 0 & 0 & 0 & \vec{E}\vec{B} \end{vmatrix} . \tag{13.14}$$

Using (13.13) and (13.14), we find the explicit form of the convolution of four tensors:

$$(F_\alpha{}^\beta)(F_\beta{}^\chi)(F_\chi{}^\sigma)(F_\sigma{}^\rho) = (\vec{E}^{\,2} - \vec{B}^{\,2})(F_\alpha{}^\beta)(F_\beta{}^\rho) + (\vec{E}\vec{B})^{\,2}\delta_\alpha{}^\rho \, . \tag{13.15}$$

Relationship (13.15) provides us with the minimal polynomial of the electromagnetic matrix $(F_\alpha{}^\beta)$. The inverse of the $\Lambda_\sigma{}^\alpha = \mu\delta_\sigma{}^\alpha + \lambda F_\sigma{}^\alpha$, the matrix $(\Lambda^{-1})_\alpha{}^\beta$, is to be searched in the form the following linear combination

$$(\Lambda^{-1})_\alpha{}^\beta = \lambda_1\, \delta_\alpha{}^\beta + \lambda_2\, F_\alpha{}^\beta + \lambda_3\, F_\alpha{}^\rho F_\rho{}^\beta + \lambda_4\, F_\alpha{}^\rho F_\rho{}^\sigma F_\sigma{}^\beta \, . \tag{13.16}$$

From the identity

$$\begin{aligned} \Lambda_\sigma{}^\alpha\, (\Lambda^{-1})_\alpha{}^\beta &= \{\mu\delta_\delta{}^\alpha + \lambda F_\sigma{}^\alpha\}\{\lambda_1\delta_\alpha{}^\beta + \lambda_2 F_\alpha{}^\beta + \lambda_3 F_\alpha{}^\rho F_\rho^\beta + \lambda_4 F_\alpha{}^\rho F_\rho{}^\chi F_\chi{}^\beta\} \\ &= \mu\lambda_1\delta_\sigma{}^\beta + \mu\lambda_2 F_\sigma{}^\beta + \mu\lambda_3 F_\sigma{}^\rho F_\rho{}^\beta + \mu\lambda_4 F_\sigma{}^\rho F_\rho{}^\chi F_\chi{}^\beta + \lambda\lambda_1\, F_\sigma{}^\beta \\ &+\lambda\lambda_2\, F_\sigma{}^\alpha F_\alpha^\beta + \lambda\lambda_3\, F_\sigma{}^\alpha F_\alpha{}^\rho F_\rho{}^\beta + \lambda\lambda_4\, F_\sigma{}^\alpha F_\alpha{}^\rho F_\rho{}^\chi F_\chi{}^\beta = \delta_\sigma{}^\beta \, . \end{aligned} \tag{13.17}$$

we get the linear non-homogeneous system of equations for parameters $\lambda_1, \lambda_2, \lambda_3, \lambda_4$:

$$\begin{aligned} \mu\, \lambda_1 + \lambda\lambda_4\, (\vec{E}\vec{B})^2 = 1 \, , \; \lambda\, \lambda_1 + \mu\, \lambda_2 = 0 \, , \\ \lambda\lambda_2 + \mu\lambda_3 + \lambda\lambda_4\, (\vec{E}^2 - \vec{B}^2) = 0 \, , \; \lambda\lambda_3 + \mu\, \lambda_4 = 0 \, . \end{aligned} \tag{13.18}$$

The determinant of the main matrix is

$$\begin{vmatrix} \mu & 0 & 0 & \lambda(\vec{E}\vec{B})^2 \\ \lambda & \mu & 0 & 0 \\ 0 & \lambda & \mu & \lambda(\vec{E}^2 - \vec{B}^2) \\ 0 & 0 & \lambda & \mu \end{vmatrix} = \mu^2 \left[\mu^2 - \lambda^2(\vec{E}^2 - \vec{B}^2)\right] - \lambda^4(\vec{E}\vec{B})^2 \, . \tag{13.19}$$

By Cramer's rule we obtain the solution of the linear system:

$$\lambda_1 = \frac{\mu[\mu^2 - \lambda^2(\vec{E}^2 - \vec{B}^2)]}{\mu^2[\mu^2 - \lambda^2(\vec{E}^2 - \vec{B}^2)] - \lambda^4(\vec{E}\vec{B})^2} ,$$
$$\lambda_2 = \frac{-\lambda[\mu^2 - \lambda^2(\vec{E}^2 - \vec{B}^2)]}{\mu^2[\mu^2 - \lambda^2(\vec{E}^2 - \vec{B}^2)] - \lambda^4(\vec{E}\vec{B})^2} ,$$
$$\lambda_3 = \frac{\mu\lambda^2}{\mu^2[\mu^2 - \lambda^2(\vec{E}^2 - \vec{B}^2)] - \lambda^4(\vec{E}\vec{B})^2} ,$$
$$\lambda_4 = -\frac{\lambda^3}{\mu^2[\mu^2 - \lambda^2(\vec{E}^2 - \vec{B}^2)] - \lambda^4(\vec{E}\vec{B})^2} . \tag{13.20}$$

Additionally, we check the validity of the following relations:

$$F_\alpha{}^\beta F_\beta{}^\alpha = 2(\vec{E}^2 - \vec{B}^2) , \qquad F_\alpha{}^\beta F_\beta{}^{\times\rho} = 4\vec{E}\vec{B}\delta_\alpha^\beta , \qquad \vec{E}\vec{B} = \frac{1}{4}F_\alpha{}^\beta F_\beta{}^{\times\rho} .$$

Thus, we arrive at the following explicit representation of the inverse matrix:

$$(\Lambda^{-1})_\alpha{}^\beta = \frac{1}{\mu^2\left(\mu^2 - \frac{\lambda^2}{2}F_\rho{}^\sigma F_\sigma{}^\rho\right) - \lambda^4\left(\frac{1}{4}F_\alpha{}^\beta F_\beta^{\times\rho}\right)^2} \tag{13.21}$$

$$\times\left\{\mu\left(\mu^2 - \frac{\lambda^2}{2}F_\rho{}^\sigma F_\sigma{}^\rho\right)\ \delta_\alpha^\beta - \lambda\left(\mu^2 - \frac{\lambda^2}{2}F_\rho{}^\sigma F_\sigma{}^\rho\right)F_\alpha{}^\beta + \mu\lambda^2\ \ F_\alpha{}^\sigma F_\sigma{}^\beta - \lambda^3\ \ F_\alpha{}^\sigma F_\sigma{}^\delta F_\delta{}^\beta\right\} .$$

Now let us consider the task of finding the inverse matrix $(\Lambda^{-1})_\alpha{}^\beta$ in the case of Riemannian space. For simplicity, we will assume the metric tensor be diagonal. It insures the following property of the electromagnetic tensor $F_{\alpha\beta}$:

$$F_{00} = 0 \Longrightarrow F_0{}^0 = 0 , \qquad F_{11} = 0 \Longrightarrow F_1{}^1 = 0 ,$$
$$F_{22} = 0 \Longrightarrow F_2{}^2 = 0 , \qquad F_{33} = 0 \Longrightarrow F_3{}^3 = 0 .$$

We start from the explicit form of the tensor $(F_\alpha{}^\beta)$:

$$(F_\alpha{}^\beta) = \begin{vmatrix} 0 & F_0{}^1 & F_0{}^2 & F_0{}^3 \\ F_1{}^0 & 0 & F_1{}^2 & F_1{}^3 \\ F_2{}^0 & F_2{}^1 & 0 & F_2{}^3 \\ F_3{}^0 & F_3{}^1 & F_3{}^2 & 0 \end{vmatrix} = \begin{vmatrix} 0 & g^{11}E_1 & g^{22}E_2 & g^{33}E_3 \\ -g^{00}E_1 & 0 & g^{22}B_3 & -g^{33}B_2 \\ -g^{00}E_2 & -g^{11}B_3 & 0 & g^{33}B_1 \\ -g^{00}E_3 & g^{11}B_2 & -g^{22}B_1 & 0 \end{vmatrix} . \tag{13.22}$$

Below we will use the notation

$$g^{11}E_1 = E^1,\ g^{22}E_2 = E^2,\ g^{33}E_3 = E^3,\ \ g^{00} = h ,$$
$$g^{22}g^{33}B_1 = B^1 ,\ g^{33}g^{11}B_2 = B^2 ,\ g^{11}g^{22}B_3 = B^3 .$$

Let us compute the convolution of two tensors (we write the matrix by the columns)

$$(F_\alpha{}^\beta)(F_\beta{}^\rho) = \begin{vmatrix} -hE^iE_i & -(E_2B^3 - E_3B^2) \\ -h(E^2B_3 - E^3B_2) & -hE_1E^1 - B_2B^2 - B_3B^3 \\ -h(E^3B_1 - E^1B_3) & -hE^1E_2 + B_1B^2 \\ -h(E^1B_2 - E^2B_1) & -hE^1E_3 + B_1B^3 \end{vmatrix} ;$$

$$\left|\begin{array}{cc} -(E_3B^1 - E_1B^3) & -(E_1B^2 - E_2B^1) \\ -hE_1E^2 + B^1B_2 & -hE_1E^3 + B^1B_3 \\ -hE_2E^2 - B_1B^1 - B_3B^3 & -hE_2E^3 + B^2B_3 \\ -hE^2E_3 + B_2B^3 & -hE_3E^3 - B_1B^1 - B_2B^2 \end{array}\right| ;$$

next, compute the convolution of three tensors

$$(F_\alpha{}^\beta)(F_\beta{}^\rho)(F_\rho{}^\sigma) = -(\, g^{00}E_iE^i + B_iB^i \,) \times \left|\begin{array}{cccc} 0 & E^1 & E^2 & E^3 \\ -g^{00}E_1 & 0 & g^{22}B_3 & -g^{33}B_2 \\ -g^{00}E_2 & -g^{11}B_3 & 0 & g^{33}B_1 \\ -g^{00}E_3 & g^{11}B_2 & -g^{22}B_1 & 0 \end{array}\right| + (B_iE_i) \left|\begin{array}{cccc} 0 & g^{11}B^1 & g^{22}B^2 & g^{33}B^3 \\ -g^{00}B^1 & 0 & g^{22}g^{00}E^3 & -g^{33}g^{00}E^2 \\ -g^{00}B^2 & -g^{11}g^{00}E^3 & 0 & g^{33}g^{00}E^1 \\ -g^{00}B^3 & g^{11}g^{00}E^2 & -g^{22}g^{00}E^1 & 0 \end{array}\right| . \tag{13.23}$$

Using (**??**), we compute the convolution of two tensors in two pairs of indices

$$\frac{1}{2}\,(F_\alpha{}^\beta F_\beta{}^\alpha) = -(\, g^{00}E_iE^i + B_iB^i \,) \equiv I(x) \,. \tag{13.24}$$

Let us specify the dual electromagnetic tensor

$$(F^\times)^{\alpha\beta} = \frac{1}{2}\epsilon^{\alpha\beta\rho\sigma}(x)F_{\rho\sigma} \,, \qquad \epsilon^{0123}(x) = \epsilon(x), \qquad \epsilon^{\alpha\beta\rho\sigma}(x) = \epsilon^{[\alpha\beta\rho\sigma]}(x) \,, \tag{13.25}$$

where

$$\epsilon(x) = \frac{1}{\sqrt{-\det g}} = \frac{1}{\sqrt{-g_{00}g_{11}g_{22}g_{33}}} = \sqrt{-g^{00}g^{11}g^{22}g^{33}} \,. \tag{13.26}$$

Further, we find explicit expressions for the components of the dual tensor:

$$(F^\times)^{01} = \epsilon^{0123}(x)F_{23} = \epsilon(x)F_{23} = \epsilon(x)B_1 \,, \qquad (F^\times)_0{}^1 = g_{00}\epsilon(x)B_1 = -\sqrt{-g}\; g^{11}B^1 \,,$$

$$(F^\times)_1{}^0 = -g_{11}\epsilon(x)B_1 = +\sqrt{-g}\; g^{00}B^1 \,, \qquad (F^\times)^{02} = \epsilon^{0231}(x)F_{31} = \epsilon(x)F_{31} = \epsilon(x)B_2 \,,$$

$$(F^\times)_0{}^2 = g_{00}\epsilon(x)B_2 = -\sqrt{-g}\; g^{22}B^2 \,, \qquad (F^\times)_2{}^0 = -g_{22}\epsilon(x)B_2 = +\sqrt{-g}\; g^{00}B^2 \,,$$

$$(F^\times)^{03} = \epsilon^{0312}(x)F_{12} = \epsilon(x)F_{12} = \epsilon(x)B_3 \,, \qquad (F^\times)_0{}^3 = g_{00}\epsilon(x)B_3 = -\sqrt{-g}\; g^{33}B^3 \,,$$

$$(F^\times)_3{}^0 = -g_{33}\epsilon(x)B_3 = +\sqrt{-g}\; g^{00}B^3 \,, \qquad (F^\times)^{23} = \epsilon^{2301}(x)F_{01} = \epsilon(x)F_{01} = \epsilon(x)E_1 \,,$$

$$(F^\times)_2{}^3 = g_{22}\epsilon(x)E_1 = -\sqrt{-g}\; g^{00}g^{33}E^1 \,, \qquad (F^\times)_3{}^2 = -g_{33}\epsilon(x)E_1 = +\sqrt{-g}\; g^{00}g^{22}E^1 \,,$$

$$(F^\times)^{31} = \epsilon^{3102}(x)F_{02} = \epsilon(x)F_{02} = \epsilon(x)E_2 \,, \qquad (F^\times)_3{}^1 = g_{33}\epsilon(x)E_2 = -\sqrt{-g}\; g^{00}g^{11}E^2 \,,$$

$$(F^\times)_1{}^3 = -g_{11}\epsilon(x)E_2 = +\sqrt{-g}\; g^{00}g^{33}E^2 \,, \qquad (F^\times)^{12} = \epsilon^{1203}(x)F_{03} = \epsilon(x)F_{03} = \epsilon(x)E_3 \,,$$

$$(F^\times)_1{}^2 = g_{11}\epsilon(x)E_3 = -\sqrt{-g}\; g^{00}g^{22}E^3 \,, \qquad (F^\times)_2{}^1 = -g_{22}\epsilon(x)E_3 = +\sqrt{-g}\; g^{00}g^{11}E^3 \,.$$

So we obtain

$$\frac{1}{4}(F_\alpha^{\times\beta})(F_\beta{}^\alpha) = \frac{1}{4}\mathrm{Sp}\left\{ -\sqrt{-g}\begin{vmatrix} 0 & g^{11}B^1 & g^{22}B^2 & g^{33}B^3 \\ -g^{00}B^1 & 0 & g^{00}g^{22}E^3 & -g^{00}g^{33}E^2 \\ -g^{00}B^2 & -g^{00}g^{11}E^3 & 0 & g^{00}g^{33}E^1 \\ -g^{00}B^3 & g^{00}g^{11}E^2 & -g^{00}g^{22}E^1 & 0 \end{vmatrix} \right.$$

$$\left. \times \begin{vmatrix} 0 & E^1 & E^2 & E^3 \\ -g^{00}E_1 & 0 & g^{22}B_3 & -g^{33}B_2 \\ -g^{00}E_2 & -g^{11}B_3 & 0 & g^{33}B_1 \\ -g^{00}E_3 & g^{11}B_2 & -g^{22}B_1 & 0 \end{vmatrix} \right\} ;$$

let us consider expressions for diagonal elements of this product:

$$(00) = \sqrt{-g}\; g^{00}g^{11}g^{22}g^{33}(E_iB_i) = -\frac{1}{\sqrt{-g}}(E_iB_i) ,$$

$$(11) = -\frac{1}{\sqrt{-g}}(E_iB_i) , \qquad (22) = -\frac{1}{\sqrt{-g}}(E_iB_i) , \qquad (33) = -\frac{1}{\sqrt{-g}}(E_iB_i) .$$

Thus we arrive at the relationship

$$\frac{1}{4}(F_\alpha^{\times\beta})(F_\beta{}^\rho) = -\frac{1}{\sqrt{-g}}\;(E_iB_i) \equiv J(x) . \tag{13.27}$$

Turning now to (13.23) we derive the following expansion for the convolution of three tensors

$$F_\alpha{}^\beta F_\beta{}^\rho F_\rho{}^\sigma = I(x)\; F_\alpha{}^\sigma + J(x)\; F_\alpha^{\times\,\sigma} ; \tag{13.28}$$

where the notation $I(x), J(x)$ for the two invariants of the electromagnetic field is used. An important point should be stressed. The representation (13.28), due to the tensor nature of the relation, is valid in all (including non-orthogonal) coordinate systems of space–time.

With the help of (13.28) we easily find the explicit form of the convolution of the four tensors

$$(F_\alpha{}^\beta F_\beta{}^\chi F_\chi{}^\sigma)(F_\sigma{}^\rho) = I(x)\; F_\alpha{}^\sigma F_\sigma{}^\rho - J(x)\;\frac{1}{\sqrt{-g}}\;(E_iB_i)\delta_\alpha{}^\rho = I(x)\; F_\alpha{}^\sigma F_\sigma{}^\rho + J^2(x)\;\delta_\alpha{}^\rho . \tag{13.29}$$

At the same time, the relation (13.29) defines the minimal polynomial of the matrix $(F_\alpha{}^\beta)$.

The inverse of $\Lambda_\sigma{}^\alpha = \mu\delta_\sigma{}^\alpha + \lambda F_\sigma{}^\alpha$ tensor $(\Lambda^{-1})_\alpha{}^\beta$ should be constructed in the form

$$(\Lambda^{-1})_\alpha{}^\beta = \lambda_1\;\delta_\alpha{}^\beta + \lambda_2\;F_\alpha{}^\beta + \lambda_3\;F_\alpha{}^\rho F_\rho{}^\beta + \lambda_4\;F_\alpha{}^\rho F_\rho{}^\sigma F_\sigma{}^\beta . \tag{13.30}$$

From the identity

$$\begin{aligned} \Lambda_\sigma{}^\alpha\;(\Lambda^{-1})_\alpha{}^\beta &= \{\mu\delta_\delta{}^\alpha + \lambda F_\sigma{}^\alpha\}\{\lambda_1\delta_\alpha{}^\beta + \lambda_2 F_\alpha{}^\beta + \lambda_3 F_\alpha{}^\rho F_\rho^\beta + \lambda_4 F_\alpha{}^\rho F_\rho{}^\chi F_\chi{}^\beta\} \\ &= \mu\lambda_1\delta_\sigma{}^\beta + \mu\lambda_2 F_\sigma{}^\beta + \mu\lambda_3 F_\sigma{}^\rho F_\rho{}^\beta + \mu\lambda_4 F_\sigma{}^\rho F_\rho{}^\chi F_\chi{}^\beta \\ &+\lambda\lambda_1\;F_\sigma{}^\beta + \lambda\lambda_2\;F_\sigma{}^\alpha F_\alpha^\beta + \lambda\lambda_3\;F_\sigma{}^\alpha F_\alpha{}^\rho F_\rho{}^\beta + \lambda\lambda_4\;F_\sigma{}^\alpha F_\alpha{}^\rho F_\rho{}^\chi F_\chi{}^\beta = \delta_\sigma{}^\beta . \end{aligned} \tag{13.31}$$

it follows the linear non-homogeneous system of equations for parameters $\lambda_1, \lambda_2, \lambda_3, \lambda_4$:

$$\mu\lambda_1 + \lambda\lambda_4 J^2 x = 1 ,\; \lambda\lambda_1 + \mu\lambda_2 = 0 ,\; \lambda\lambda_2 + \mu\lambda_3 + \lambda\lambda_4\; I(x) = 0 ,\; \lambda\lambda_3 + \mu\lambda_4 = 0 ; \tag{13.32}$$

it solution is

$$\lambda_1 = \frac{\mu(\mu^2 - \lambda^2 I)}{\mu^2(\mu^2 - \lambda^2 I) - \lambda^4 J^2}\,, \qquad \lambda_2 = \frac{-\lambda(\mu^2 - \lambda^2 I)}{\mu^2(\mu^2 - \lambda^2 I) - \lambda^4 J^2}\,,$$
$$\lambda_3 = \frac{\mu\lambda^2}{\mu^2(\mu^2 - \lambda^2 I) - \lambda^4 J^2}\,, \qquad \lambda_4 = -\frac{\lambda^3}{\mu^2(\mu^2 - \lambda^2 I) - \lambda^4 J^2}\,. \tag{13.33}$$

Thus, we arrive at the following explicit representation of the inverse tensor:

$$(\Lambda^{-1})_\alpha{}^\beta = \frac{1}{\mu^2\,(\mu^2 - \lambda^2 I) - \lambda^4 J^2}$$
$$\times \Big\{\mu\,(\,\mu^2 - \lambda^2 I\,)\,\delta_\alpha^\beta - \lambda\,(\mu^2 - \lambda^2 I)\,F_\alpha{}^\beta + \mu\lambda^2\,F_\alpha{}^\sigma F_\sigma{}^\beta - \lambda^3\;F_\alpha{}^\sigma F_\sigma{}^\delta F_\delta{}^\beta\Big\}\,. \tag{13.34}$$

Using (13.28), $F_\alpha{}^\sigma F_\sigma{}^\delta F_\delta{}^\beta = I\,F_\alpha{}^\beta + J\,F_\alpha^{\times\,\beta}$, we find a more simple representation

$$(\Lambda^{-1})_\alpha{}^\beta = \frac{1}{\mu^2(\mu^2 - \lambda^2 I) - \lambda^4 J^2}\Big\{\mu\,(\,\mu^2 - \lambda^2 I\,)\,\delta_\alpha^\beta - \lambda\mu^2\,F_\alpha{}^\beta + \mu\lambda^2\,F_\alpha{}^\sigma F_\sigma{}^\beta - \lambda^3\,J(x)\,F_\alpha^{\times\beta}\Big\}\,. \tag{13.35}$$

These results will be significantly simplified in the case of a purely magnetic or purely electric field.

Magnetic field:

$$(\Lambda^{-1})_\alpha{}^\beta = \frac{1}{\mu^2(\mu^2 - \lambda^2 I)}\Big\{\mu\,(\,\mu^2 - \lambda^2 I\,)\,\delta_\alpha^\beta - \lambda\mu^2\,F_\alpha{}^\beta + \mu\lambda^2\,F_\alpha{}^\sigma F_\sigma{}^\beta\Big\}\,,$$
$$I(x) = -(B_i B^i\,)\,, \qquad J(x) = 0\,, \qquad (F_\alpha{}^\beta) = \begin{vmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & g^{22}B_3 & -g^{33}B_2 \\ 0 & -g^{11}B_3 & 0 & g^{33}B_1 \\ 0 & g^{11}B_2 & -g^{22}B_1 & 0 \end{vmatrix},$$
$$(F_\alpha{}^\sigma)(F_\sigma{}^\beta) = \begin{vmatrix} 0 & 0 & 0 & 0 \\ 0 & -B_2B^2 - B_3B^3 & B^1B_2 & B^1B_3 \\ 0 & B_1B^2 & -B_1B^1 - B_3B^3 & B^2B_3 \\ 0 & B_1B^3 & B_2B^3 & -B_1B^1 - B_2B^2 \end{vmatrix}. \tag{13.36}$$

Electric field:

$$(\Lambda^{-1})_\alpha{}^\beta = \frac{1}{\mu^2(\mu^2 - \lambda^2 I)}\Big\{\mu\,(\,\mu^2 - \lambda^2 I)\,\delta_\alpha^\beta - \lambda\mu^2\,F_\alpha{}^\beta + \mu\lambda^2\,F_\alpha{}^\sigma F_\sigma{}^\beta\Big\}\,,$$
$$I = -(\,g^{00}E_iE^i)\,, \qquad J = 0\,, \qquad (F_\alpha{}^\beta) = \begin{vmatrix} 0 & E^1 & E^2 & E^3 \\ -g^{00}E_1 & 0 & 0 & 0 \\ -g^{00}E_2 & 0 & 0 & 0 \\ -g^{00}E_3 & 0 & 0 & 0 \end{vmatrix},$$
$$(F_\alpha{}^\sigma)(F_\sigma{}^\beta) = g^{00}\begin{vmatrix} -E^iE_i & 0 & 0 & 0 \\ 0 & -E_1E^1 & -E_1E^2 & -E_1E^3 \\ 0 & -E^1E_2 & -E_2E^2 & -E_2E^3 \\ 0 & -E^1E_3 & -E^2E_3 & -E_3E^3- \end{vmatrix}. \tag{13.37}$$

Now we can construct in explicit form the generalized scalar equation, the tensor notation. Substituting the above tensor $(\Lambda^{-1})_\alpha{}^\beta$ into the equation (13.4) for a scalar function we obtain

$$\Big\{\frac{\mu}{\det\Lambda}\Big(\mu\,(\,\mu^2-\lambda^2 I\,)\,\delta_\alpha^\beta-\lambda\mu^2\,F_\alpha{}^\beta+\mu\lambda^2\,F_\alpha{}^\sigma F_\sigma{}^\beta-\lambda^3\,J\,F_\alpha^{\times\beta}\Big)\,D^\alpha D_\beta+ \tag{13.38}$$

$$+\Big[i\hbar\mu\partial^\alpha\frac{1}{\det\Lambda}\Big(\mu\,(\,\mu^2-\lambda^2 I\,)\,\delta_\alpha^\beta-\lambda\mu^2\,F_\alpha{}^\beta+\mu\lambda^2\,F_\alpha{}^\sigma F_\sigma{}^\beta-\lambda^3\,J\,F_\alpha^{\times\beta}\Big)\Big]\,D_\beta-\mu^2\Big\}\,\Phi=0\,.$$

In the case of constant and uniform fields (in Cartesian coordinates of the flat space), eq. (13.38) becomes much more simple:

$$\Big\{\mu\Big[\mu\,(\,\mu^2-\lambda^2 I\,)\,\delta_\alpha^\beta-\lambda\mu^2\,F_\alpha{}^\beta+\mu\lambda^2\,F_\alpha{}^\sigma F_\sigma{}^\beta-\lambda^3\,J\,F_\alpha^{\times\beta}\Big]\,D^\alpha D_\beta-\mu^2\det\Lambda\Big\}\,\Phi=0\,. \tag{13.39}$$

The generalization of (13.38) to the case of space-time models with the Riemannian geometry is achieved by a formal change of ordinary derivatives by covariant derivatives: $\partial_\alpha\longrightarrow\nabla_\alpha$, $D_\alpha=i\hbar\nabla_\alpha-\frac{e}{c}A_\alpha$, as a result we have

$$\begin{aligned}\mu\Big[\mu\,(\,\mu^2-\lambda^2 I\,)\,\delta_\alpha^\beta-\lambda\mu^2\,F_\alpha{}^\beta+\mu\lambda^2\,F_\alpha{}^\sigma F_\sigma{}^\beta-\lambda^3\,J\,F_\alpha^{\times\beta}\Big]\,D^\alpha D_\beta\Phi\\ +i\hbar\mu\,\det\Lambda\Big[\nabla_\alpha\Big(\frac{\mu\,(\,\mu^2-\lambda^2 I}{\det\Lambda}\,)\,g^{\alpha\beta}-\frac{\lambda\mu^2}{\det\Lambda}\,F^{\alpha\beta}\\ -\frac{\lambda^3\,J}{\det\Lambda}\,F^{\times\alpha\beta}+\frac{\mu\lambda^2}{\det\Lambda}\,g_{\rho\sigma}F^{\alpha\rho}F^{\sigma\beta}\Big)\Big]\,D_\beta\Phi-\mu^2\det\Lambda\,\Phi\;=0\,.\end{aligned} \tag{13.40}$$

Let us specify the separate terms under the symbol of the covariant derivative in

$$\Big[\nabla_\alpha\Big(\frac{\mu\,(\,\mu^2-\lambda^2 I}{\det\Lambda}\,)\,g^{\alpha\beta}-\frac{\lambda\mu^2}{\det\Lambda}\,F^{\alpha\beta}-\frac{\lambda^3\,J}{\det\Lambda}\,F^{\times\alpha\beta}+\frac{\mu\lambda^2}{\det\Lambda}\,g_{\rho\sigma}F^{\alpha\rho}F^{\sigma\beta}\Big)\Big]\,D_\beta\Phi\,.$$

The first term is reduced to the ordinary derivative of a scalar. The second and third terms are the 4-divergences of the antisymmetric tensor, and the fourth term is a 4-divergence of the symmetric tensor; they are calculated using known formulas

$$\nabla_\alpha(A^{\alpha\beta})=(\frac{1}{\sqrt{-g}}\partial_\alpha\sqrt{-g}A^{\alpha\beta})\,,\qquad \nabla_\alpha(S^\alpha{}_\beta)=(\frac{1}{\sqrt{-g}}\partial_\alpha\sqrt{-g}S^\alpha{}_\beta)-\frac{1}{2}(\partial_\beta g_{\rho\sigma})S^{\rho\sigma}\,. \tag{13.41}$$

Now, let us derive non-relativistic wave equation for Cox's particle, stating with

$$K_\rho{}^\alpha\Big(i\,\nabla_\alpha-\frac{e}{c\hbar}A_\alpha\Big)\;\Phi=\frac{Mc}{\hbar}\Phi_\rho\,,\qquad\Big(i\,\nabla_\alpha-\frac{e}{c\hbar}A_\alpha\Big)\;\Phi^\alpha=\frac{Mc}{\hbar}\Phi\,. \tag{13.42}$$

The equations (13.42) can be rewritten in a form more convenient for practical calculations:

$$K_\rho{}^\alpha(i\,\partial_\alpha-\frac{e}{c\hbar}A_\alpha)\Phi=\frac{Mc}{\hbar}\Phi_\rho\,,\qquad(\frac{i}{\sqrt{-g}}\,\frac{\partial}{\partial x^\alpha}\sqrt{-g}-\frac{e}{c\hbar}\,A_\alpha)\,g^{\alpha\beta}\Phi_\beta=\frac{Mc}{\hbar}\Phi\,. \tag{13.43}$$

Considering the space-time models with the metric $dS^2=c^2dt^2+g_{kl}(x)\,dx^kdx^l$, in (13.43) let us perform the splitting (3 +1)

$$\begin{aligned}K_0{}^0(i\,\partial_0-\frac{e}{c\hbar}A_0)\,\Phi+K_0{}^k(i\,\partial_k-\frac{e}{c\hbar}A_k)\,\Phi=\frac{Mc}{\hbar}\,\Phi_0\,,\\ K_j{}^0(i\,\partial_0-\frac{e}{c\hbar}A_0)\,\Phi+K_j{}^k(i\,\partial_k-\frac{e}{c\hbar}A_k)\,\Phi=\frac{Mc}{\hbar}\,\Phi_j\,,\\ (\frac{i}{\sqrt{-g}}\,\frac{\partial}{\partial x^0}\sqrt{-g}-\frac{e}{c\hbar}\,A_0)\,\Phi_0+(\frac{i}{\sqrt{-g}}\,\frac{\partial}{\partial x^k}\sqrt{-g}-\frac{e}{c\hbar}\,A_k)\,g^{kl}\Phi_l=\frac{Mc}{\hbar}\Phi\,.\end{aligned} \tag{13.44}$$

Next, separate the rest energy by the substitutions

$$\Phi \Longrightarrow \exp\,(-i\frac{Mc^2t}{\hbar})\Phi\ , \qquad \Phi_0 \Longrightarrow \exp\,(-i\frac{Mc^2t}{\hbar})\Phi_0\ , \qquad \Phi_l \Longrightarrow \exp\,(-i\frac{Mc^2t}{\hbar})\Phi_l\ .$$

As a result, from of (13.44) we get

$$K_0{}^0(i\hbar\,\partial_t + Mc^2 - eA_0)\,\Phi + K_0{}^k(ic\hbar\,\partial_k - eA_k)\,\Phi = Mc^2\,\Phi_0\ , \tag{13.45}$$

$$K_j{}^0(i\hbar\,\partial_t + Mc^2 - eA_0)\,\Phi + K_j{}^k(ic\hbar\,\partial_k - eA_k)\,\Phi = Mc^2\,\Phi_j\ , \tag{13.46}$$

$$\left(i\hbar\partial_t + Mc^2 + \frac{i\hbar}{\sqrt{-g}}\,\frac{\partial\sqrt{-g}}{\partial t} - e\,A_0\right)\,\Phi_0 + \left(\frac{ic\hbar}{\sqrt{-g}}\,\frac{\partial}{\partial x^k}\sqrt{-g} - e\,A_k\right)\,g^{kj}\Phi_j = Mc^2\Phi\ . \tag{13.47}$$

Using (13.46), now exclude the vector (non-dynamic) variable Φ_j:

$$K_0{}^0(i\hbar\,\partial_t + Mc^2 - eA_0)\,\Phi + K_0{}^k(ic\hbar\,\partial_k - eA_k)\,\Phi = Mc^2\,\Phi_0\ , \tag{13.48}$$

$$\left(i\hbar\partial_t + Mc^2 + \frac{i\hbar}{\sqrt{-g}}\,\frac{\partial\sqrt{-g}}{\partial t} - e\,A_0\right)\,\Phi_0 + \left(\frac{ic\hbar}{\sqrt{-g}}\,\frac{\partial}{\partial x^k}\sqrt{-g} - e\,A_k\right)$$
$$\times\frac{g^{kj}}{mc^2}\left[K_j{}^0(i\hbar\,\partial_t + Mc^2 - eA_0) + K_j{}^l(ic\hbar\,\partial_l - eA_l)\right]\Phi = Mc^2\Phi\ . \tag{13.49}$$

With notation

$$i\hbar\,\partial_t - eA_0 = D_t\ , \qquad ic\hbar\,\partial_k - eA_k = cD_k\ , \qquad \frac{ic\hbar}{\sqrt{-g}}\,\frac{\partial}{\partial x^k}\sqrt{-g} - e\,A_k = c\,\overset{\circ}{D}_k\ , \tag{13.50}$$

equations (13.48), (13.49) can be written as follows

$$K_0{}^0(D_t + Mc^2)\,\Phi + K_0{}^k cD_k\,\Phi = Mc^2\Phi_0\ ,$$

$$\left(D_t + Mc^2 + \frac{i\hbar}{\sqrt{-g}}\,\frac{\partial\sqrt{-g}}{\partial t}\right)\Phi_0 + \overset{\circ}{D}_k\ \frac{g^{kj}}{M}\left(\frac{1}{c}K_j{}^0(D_t + Mc^2) + K_j{}^l D_l\right)\Phi = Mc^2\Phi\ . \tag{13.51}$$

Following the method described in [3], we introduce the small component φ and the big component Ψ: $\Phi = (\Psi + \varphi)/2,\ \Phi_0 = (\Psi - \varphi)/2$. Substituting these relations in (13.51):

$$K_0{}^0 D_t\frac{\Psi+\varphi}{2} + (K_0{}^0 - 1 + 1)Mc^2\,\frac{\Psi+\varphi}{2} + K_0{}^k cD_k\,\frac{\Psi+\varphi}{2} = Mc^2\,\frac{\Psi-\varphi}{2}\ ,$$

$$\left(D_t + Mc^2 + \frac{i\hbar}{\sqrt{-g}}\,\frac{\partial\sqrt{-g}}{\partial t}\right)\,\frac{\Psi-\varphi}{2} + \overset{\circ}{D}_k\,\frac{g^{kj}}{M}\left(\frac{1}{c}K_j{}^0(D_t + Mc^2) + K_j{}^l D_l\right)\frac{\Psi+\varphi}{2} = Mc^2\frac{\Psi+\varphi}{2}\ .$$

From whence, neglecting the small component φ in comparison with the big Ψ, we obtain

$$\left(D_t + (K_0{}^0 - 1)(D_t + Mc^2) + K_0{}^k cD_k\right)\frac{\Psi}{2} = -Mc^2\,\varphi\ , \tag{13.52}$$

$$\left(D_t + \frac{i\hbar}{\sqrt{-g}}\,\frac{\partial\sqrt{-g}}{\partial t}\right)\,\frac{\Psi}{2} + \overset{\circ}{D}_k\,\frac{g^{kj}}{M}\left(\frac{1}{c}K_j{}^0(D_t + Mc^2) + K_j{}^l D_l\right)\frac{\Psi}{2} = Mc^2\varphi\ . \tag{13.53}$$

We assume that the energy of nonrelativistic particles are much smaller than the rest energy, that is, we apply the approximation $(D_t + Mc^2) \approx Mc^2$. As a result, the first and second equations are simplified

$$\left(D_t + (K_0{}^0 - 1)Mc^2 + K_0{}^k cD_k\right)\frac{\Psi}{2} = -Mc^2\,\varphi\ , \tag{13.54}$$

$$\left(D_t + \frac{i\hbar}{\sqrt{-g}}\,\frac{\partial\sqrt{-g}}{\partial t}\right)\frac{\Psi}{2} + \overset{\circ}{D}_k\,\frac{g^{kj}}{M}\left(McK_j{}^0 + K_j{}^l D_l\right)\frac{\Psi}{2} = Mc^2\varphi\ . \tag{13.55}$$

With the help of (13.54), one can eliminate the small component φ from the equation (13.55), so we get

$$(D_t + \frac{i\hbar}{\sqrt{-g}}\,\frac{\partial}{\partial t}\sqrt{-g})\,\frac{\Psi}{2} + \overset{\circ}{D}_k\,\frac{g^{kj}}{M}\left(Mc{K_j}^0 + {K_j}^k D_k\right)\frac{\Psi}{2} = -\left(D_t + ({K_0}^0 - 1)Mc^2 + {K_0}^k c D_k\right)\frac{\Psi}{2}\,,$$

or

$$\left(D_t + \frac{i\hbar}{2\sqrt{-g}}\,\frac{\partial\sqrt{-g}}{\partial t} + \frac{1}{2}[({K_0}^0 - 1)Mc^2 + {K_0}^k c D_k]\right)\Psi = \overset{\circ}{D}_k\,\frac{(-g^{kj})}{2M}\left[{K_j}^l D_l + Mc{K_j}^0\right]\Psi\,. \quad (13.56)$$

With the substitution $\Psi \implies (-g)^{-1/4}\,\Psi$, equation (13.56) reads simpler

$$D_t\,\Psi = \frac{1}{2M}\,\overset{\circ}{D}_k\,(-g^{kj})\left({K_j}^l D_l + Mc{K_j}^0\right)\Psi - \frac{1}{2}\left(({K_0}^0 - 1)Mc^2 + {K_0}^j c D_j\right)\Psi\,. \quad (13.57)$$

This is the nonrelativistic Schrödinger equation for the Cox's particle.

Let us specify the case of the presence of only magnetic field, then the Schrödinger equation (13.57) becomes much simpler

$$D_t\,\Psi = \frac{1}{2M}\,\overset{\circ}{D}_k\,(-g^{kj}){K_j}^l D_l\,\Psi\,. \quad (13.58)$$

Let us detail the operator ${K_j}^l D_l$:

$$\begin{aligned}
{K_1}^l D_l &= {K_1}^1 D_1 + {K_1}^2 D_2 + {K_1}^3 D_3 = \frac{1}{\mu^2 + \lambda^2 B_i B^i}[\,\mu^2 D_1 + \mu\lambda(B_2 D^3 - B_3 D^2) + \lambda^2 B^1\,(B_i D_i)\,]\,,\\
{K_2}^l D_l &= {K_2}^1 D_1 + {K_2}^2 D_2 + {K_2}^3 D_3 = \frac{1}{\mu^2 + \lambda^2 B_i B^i}[\,\mu^2 D_2 + \mu\lambda(B_3 D^1 - B_1 D^3) + \lambda^2 B^2\,(\vec{B}\vec{D})\,]\,,\\
{K_3}^l D_l &= {K_3}^1 D_1 + {K_3}^2 D_2 + {K_3}^3 D_3 = \frac{1}{\mu^2 + \lambda^2 B_i B^i}[\,\mu^2 D_3 + \mu\lambda(B_1 D^2 - B_2 D^1) + \lambda^2 B^3\,(B_i D_i)\,]\,.
\end{aligned}$$

Thus, we have (we use the notation $\Gamma = \lambda/\mu$)

$$\begin{aligned}
\overset{*}{D}_1 &= {K_1}^l D_l = \frac{1}{1 + \Gamma^2 B_i B^i}\left[\,D_1 + \Gamma(B_2 D^3 - B_3 D^2) + \Gamma^2 B^1\,(B_i D_i)\,\right]\,,\\
\overset{*}{D}_2 &= {K_2}^l D_l = \frac{1}{1 + \Gamma^2 B_i B^i}\left[\,D_2 + \Gamma(B_3 D^1 - B_1 D^3) + \Gamma^2 B^2\,(\vec{B}\vec{D})\,\right]\,,\\
\overset{*}{D}_3 &= {K_3}^l D_l = \frac{1}{1 + \Gamma^2 B_i B^i}\left[\,D_3 + \Gamma(B_1 D^2 - B_2 D^1) + \Gamma^2 B^3\,(B_i D_i)\,\right]\,.
\end{aligned} \quad (13.59)$$

Therefore, the equation (13.58) can be written in a compact form as follows

$$D_t\,\Psi = -\frac{1}{2M}\,\overset{\circ}{D}_k\,g^{kj}(x)\,\overset{*}{D}_j\,\Psi\,. \quad (13.60)$$

In the case of the Cartesian coordinates of the flat space, metric tensor is trivial, and the equation is simplified

$$D_t\,\Psi = \frac{1}{2M}\vec{D}\left(\frac{\vec{D} - \Gamma\,\vec{B}\times\vec{D} + \Gamma^2\vec{B}(\vec{B}\vec{D})}{1 + \Gamma^2\vec{B}^{\,2}}\right)\Psi\,. \quad (13.61)$$

If the magnetic field in the flat space is uniform, equation (13.61) can be simplified yet more

$$D_t\,\Psi = \frac{1}{2M}\frac{1}{1+\Gamma^2\vec{B}^{\,2}}\left(\vec{D}^2 - \Gamma\;\vec{D}(\vec{B}\times\vec{D}) + \Gamma^2\;(\vec{B}\vec{D})^2\right)\Psi\;. \tag{13.62}$$

Due to the identity $\vec{D}(\vec{B}\times\vec{D}) = +i\frac{e\hbar}{c}\vec{B}^{\,2}$, the equation (13.62) can be represented as follows:

$$D_t\,\Psi = \frac{1}{2M}\frac{1}{1+\Gamma^2\vec{B}^{\,2}}\left(\vec{D}^2 - i\frac{e\hbar}{c}\Gamma\;\vec{B}^{\,2} + \Gamma^2\;(\vec{B}\vec{D})^2\right)\Psi\;. \tag{13.63}$$

Note that the presence of the term $i(e\hbar/c)\Gamma\;\vec{B}^{\,2}$ means that the parameter Γ must be purely imaginary. The explicit form of (13.63) also implies that there is a steady shift of all levels on the value determined by the amplitude of the magnetic field and the parameter $i\Gamma$.

Now consider the case of a uniform electric field.The operator ${K_j}^l D_l + \mu {K_j}^0$ is

$$\begin{aligned}
{K_1}^l D_l + \mu {K_1}^0 &= \frac{1}{1+\Gamma^2 E_i E^i}\left[\; D_1 + \Gamma^2(E_i E^i)D_1 + \Gamma^2 E_1(E^i D_i) + \mu\Gamma E_1)\;\right]\;,\\
{K_2}^l D_l + \mu {K_2}^0 &= \frac{1}{1+\Gamma^2 E_i E^i}\left[\; D_2 + \Gamma^2(E_i E^i)D_2 + \Gamma^2 E_2(E^i D_i) + \mu\Gamma E_2)\;\right]\;,\\
{K_3}^l D_l + \mu {K_3}^0 &= \frac{1}{1+\Gamma^2 E_i E^i}\left[\; D_3 + \Gamma^2(E_i E^i)D_3 + \Gamma^2 E_3(E^i D_i) + \mu\Gamma E_3)\;\right]\;,
\end{aligned}$$

thus

$$({K_j}^l D_l + \mu {K_j}^0) = \frac{1}{1+\Gamma^2 E_i E^i}\left[\; D_j + \Gamma^2(E_i E^i)D_j + \Gamma^2 E_j(E^i D_i) + \mu\Gamma E_j\;\right]. \tag{13.64}$$

So we have the representation

$$({K_0}^0 - 1)Mc^2 + {K_0}^j c D_j = -c\;\frac{\Gamma^2 E_i E^i\mu + \Gamma E^j D_j}{1+\Gamma^2 E_i E^i}\;. \tag{13.65}$$

Thus, we get

$$\left(D_t - c\;\frac{\Gamma^2 E_i E^i\mu + \Gamma E^j D_j}{2(1+\Gamma^2 E_i E^i)}\right)\Psi = \frac{1}{2M}\;\overset{\circ}{D}_k\;(-g^{kj})\left[\; D_j + \frac{\Gamma^2 E_j(E^i D_i) + \mu\Gamma E_j}{1+\Gamma^2 E_i E^i}\;\right]\Psi\,; \tag{13.66}$$

this is the Schrödinger equation for a Cox's particle in the electric field.

No we examine the initial complete Cox's system of equations [1] which includes symmetric and antisymmetric tensors:

$$\begin{aligned}
&\lambda_1 D^\beta\Phi_\beta - \mu\Phi = 0, \qquad \lambda_1^* D_\beta\Phi + \lambda_2 D^\alpha\Phi_{[\alpha\beta]} - \lambda_3 D^\alpha\Phi_{(\alpha\beta)} - \mu\Phi_\beta = 0,\\
&\lambda_2^*(D_\alpha\Phi_\beta - D_\beta\Phi_\alpha) - \mu\Phi_{[\alpha\beta]} = 0, \qquad \lambda_3^*(D_\alpha\Phi_\beta + D_\beta\Phi_\alpha - \frac{1}{2}g_{\alpha\beta}D^\rho\Phi_\rho) - \mu\Phi_{(\alpha\beta)} = 0,
\end{aligned} \tag{13.67}$$

where the auxiliary numerical parameters $\lambda_1, \lambda_2, \lambda_3$ subject to the additional constraints

$$\lambda_2\lambda_2^* - \lambda_3\lambda_3^* = 0\;,\;\; \lambda_1\lambda_1^* - \frac{3}{2}\lambda_3\lambda_3^* = 1\;; \tag{13.68}$$

symbol D_α denotes the derivative, which takes into account the presence of external electromagnetic and gravitational fields

$$D_\alpha = i\hbar\nabla_\alpha - \frac{e}{c}A_\alpha\;, \qquad \mu = Mc\;.$$

With the help of the third and the fourth equations in (13.67), let us exclude tensor components

$$\mu^{-1}\left(\lambda_2^*(D_\alpha\Phi_\beta - D_\beta\Phi_\alpha)\right) = \Phi_{[\alpha\beta]},$$
$$\mu^{-1}\lambda_3^*(D_\alpha\Phi_\beta + D_\beta\Phi_\alpha - \frac{1}{2}g_{\alpha\beta}D^\rho\Phi_\rho) = \Phi_{(\alpha\beta)}, \tag{13.69}$$

in the two other:

$$\lambda_1 D^\beta\Phi_\beta - \mu\Phi = 0\ , \tag{13.70}$$

$$\lambda_1^* D_\beta\Phi + \lambda_2 D^\alpha\mu^{-1}\left[\lambda_2^*(D_\alpha\Phi_\beta - D_\beta\Phi_\alpha)\right]$$
$$-\lambda_3 D^\alpha\mu^{-1}\lambda_3^*(D_\alpha\Phi_\beta + D_\beta\Phi_\alpha - \frac{1}{2}g_{\alpha\beta}D^\rho\Phi_\rho) - \mu\Phi_\beta = 0\ . \tag{13.71}$$

Eq. (13.71) can be presented as

$$\lambda_1^* D_\beta\Phi - \mu^{-1}(\lambda_2\lambda_2^* + \lambda_3\lambda_3^*)\ D^\alpha D_\beta\Phi_\alpha + \frac{1}{2}\mu^{-1}\lambda_3\lambda_3^*\ \ D_\beta D^\rho\Phi_\rho - \mu\Phi_\beta = 0\,. \tag{13.72}$$

In view of (13.68), one can apply identity

$$(\lambda_2\lambda_2^* + \lambda_3\lambda_3^*) = 2\lambda_3\lambda_3^*\ ;$$

hence

$$\lambda_1^* D_\beta\Phi - \mu^{-1}2\lambda_3\lambda_3^*\ D_\alpha D_\beta\Phi^\alpha + \frac{1}{2}\mu^{-1}\lambda_3\lambda_3^*\ D_\beta D_\alpha\Phi^\alpha - \mu\Phi_\beta = 0\,. \tag{13.73}$$

We use the identity

$$D_\alpha D_\beta\Phi^\alpha = D_\beta D_\alpha\Phi^\alpha + (D_\alpha D_\beta - D_\beta D_\alpha)\Phi^\alpha = D_\beta D_\alpha\Phi^\alpha + \hbar^2\left(-i\frac{e}{\hbar c}F_{\alpha\beta} - R_{\alpha\beta}\right)\Phi^{a\alpha}, \tag{13.74}$$

equation (13.73) can be converted to the following one

$$\lambda_1^* D_\beta\Phi + \mu^{-1}2\lambda_3\lambda_3^*\ \hbar^2\left(i\frac{e}{c}F_{\alpha\beta} + R_{\alpha\beta}\right)\Phi^\alpha - \frac{3}{2}\mu^{-1}\lambda_3\lambda_3^*\ D_\beta(D_\alpha\Phi^\alpha) - \mu\Phi_\beta = 0\ . \tag{13.75}$$

Taking into account the equation (13.70)

$$D_\alpha\Phi^\alpha = \frac{\mu}{\lambda_1}\Phi\ ,$$

one produces

$$\lambda_1\lambda_1^* D_\beta\Phi + \mu^{-1}2\lambda_3\lambda_3^*\ \hbar^2\left(i\frac{e}{\hbar c}F_{\alpha\beta} + R_{\alpha\beta}\right)\lambda_1\Phi^\alpha - \frac{3}{2}\lambda_3\lambda_3^*\ D_\beta\Phi - \mu\lambda_1\Phi_\beta = 0\ . \tag{13.76}$$

With the use of the second condition in (13.68)

$$\lambda_1\lambda_1^* - \frac{3}{2}\lambda_3\lambda_3^* = 1$$

on simplifies eq. (13.76) to the form

$$D_\beta\Phi + \mu^{-1}2\lambda_3\lambda_3^*\,\hbar^2\left(i\frac{e}{\hbar c}F_{\alpha\beta} + R_{\alpha\beta}\right)\lambda_1\Phi^\alpha - \mu\lambda_1\Phi_\beta = 0\,. \tag{13.77}$$

One should remember on additional equation (13.70)

$$\lambda_1 D^\beta\Phi_\beta - \mu\Phi = 0\,. \tag{13.78}$$

The parameter λ_1 can be included in the designation of vector components $\lambda_1\Phi_\beta \longrightarrow \Phi_\beta$; so we arrive at the extended Proca equations

$$D^\beta\Phi_\beta - \mu\Phi = 0\,, \quad D_\beta\Phi - \mu\Phi_\beta - i\frac{\hbar^2}{Mc}(2\lambda_3\lambda_3^*)\left(\frac{e}{\hbar c}F_{\beta\alpha} + iR_{\beta\alpha}\right)\Phi^\alpha = 0\,. \tag{13.79}$$

These equations should be compared with Proca-like system used above

$$D^\beta\Phi_\beta - \mu\Phi = 0\,, \quad D_\beta\Phi - \mu\Phi_\beta - \lambda F_{\beta\alpha}\Phi^\alpha = 0\,; \tag{13.80}$$

they correlate if (note that λ is an imaginary number)

$$\lambda = \frac{\hbar^2}{Mc}\frac{e}{\hbar c}(2i\lambda_3\lambda_3^*)\,. \tag{13.81}$$

Obviously, the system (13.79) is more general than (13.80), it takes into account non-minimal interaction of the Cox's scalar particle with external geometric background through the Ricci tensor.

Equations (13.79) can be rewritten as

$$D^\beta\Phi_\beta - \mu\Phi = 0\,, \quad D_\beta\Phi - \lambda\left(F_{\beta\alpha} + i\frac{\hbar c}{e}R_{\beta\alpha}\right)\Phi^\alpha - \mu\Phi_\beta = 0\,. \tag{13.82}$$

In the absence of the electromagnetic field, equations (13.82) are simplified (parameter $i\lambda$ is a real-valued)

$$D^\beta\Phi_\beta = \mu\Phi\,, \quad D_\beta\Phi = \left(i\lambda\frac{\hbar c}{e}R_{\beta\alpha}(x) + \mu\, g_{\beta\alpha}(x)\right)\Phi^\alpha\,. \tag{13.83}$$

This is purely geometric modification of the theory of a scalar particle in the Cox's approach.

Finally, let us find the tensor $(\Lambda^{-1})_\alpha{}^\beta$ when the Ricci tensor is taken into account. To this end, we write equation (13.83) in the form ($\lambda^* = -\lambda$; temporarily the coefficient $\frac{\hbar c}{e}$ will be a part of designation of the Ricci tensor)

$$D^\beta\Phi_\beta = \mu\Phi\,, \quad [\mu\delta_\alpha^{\ \beta} + \lambda(F_\alpha{}^\beta + iR_\alpha{}^\beta)]\Phi_\beta = D_\alpha\Phi\,. \tag{13.84}$$

With the use of the notation

$$\Lambda_\alpha{}^\beta = \mu\delta_\alpha^{\ \beta} + \lambda(F_\alpha{}^\beta + iR_\alpha{}^\beta) \tag{13.85}$$

equation (13.84) can be written as:

$$\Phi_\rho = (\Lambda^{-1})_\rho{}^\alpha D_\alpha\Phi\,, \qquad D^\rho\Phi_\rho = \mu\Phi\,. \tag{13.86}$$

From this it follows a generalized scalar equation

$$\left[D^\rho(\Lambda^{-1})_\rho{}^\alpha(x)D_\alpha \ - \ \mu\right]\Phi(x) = 0 \ . \tag{13.87}$$

Since the characteristic equation for the matrix $F_\alpha{}^\beta + iR_\alpha{}^\beta = G_\alpha{}^\beta$:

$$G^4 = g_0 + g_1 G + g_2 G^2 + g_3 G^3 \tag{13.88}$$

allows us to express the fourth power of the matrix G through I, G, G^2, G^3, we can look for the inverse matrix in the form

$$(\Lambda^{-1})_\rho{}^\alpha = \lambda_0 + \lambda_1 G + \lambda_2 G^2 + \lambda_3 G^3 \ . \tag{13.89}$$

From the equation $\Lambda\Lambda^{-1} = I$ if follows

$$\begin{aligned} I = (\mu + \lambda G)(\lambda_0 + \lambda_1 G + \lambda_2 G^2 + \lambda_3 G^3) \\ = \mu\lambda_0 + \mu\lambda_1 G + \mu\lambda_2 G^2 + \mu\lambda_3 G^3 \\ + \lambda\lambda_0 G + \lambda\lambda_1 G^2 + \lambda\lambda_2 G^3 \\ + \lambda\lambda_3(g_0 + g_1 G + g_2 G^2 + g_3 G^3) \ ; \end{aligned}$$

so we obtain the linear system

$$\begin{aligned} I: \qquad \mu\lambda_0 + \lambda\lambda_3 g_0 = 1 \ , \\ G: \qquad \mu\lambda_1 + \lambda\lambda_0 + \lambda\lambda_3 g_1 = 0 \ , \\ G^2: \qquad \mu\lambda_2 + \lambda\lambda_1 + \lambda\lambda_3 g_2 = 0 \ , \\ G^3: \qquad \mu\lambda_3 + \lambda\lambda_2 + \lambda\lambda_3 g_3 = 0 \ . \end{aligned} \tag{13.90}$$

We write the system in the matrix form

$$\begin{pmatrix} \mu & 0 & 0 & \lambda g_0 \\ \lambda & \mu & 0 & \lambda g_1 \\ 0 & \lambda & \mu & \lambda g_2 \\ 0 & 0 & \lambda & \mu + \lambda g_3 \end{pmatrix} \begin{vmatrix} \lambda_0 \\ \lambda_1 \\ \lambda_2 \\ \lambda_3 \end{vmatrix} = \begin{vmatrix} 1 \\ 0 \\ 0 \\ 0 \end{vmatrix} . \tag{13.91}$$

Its solution is

$$\begin{aligned} \lambda_0 &= \frac{-(\mu^3 + \mu^2\lambda g_3 - \mu\lambda^2 g_2 + \lambda^3 g_1)}{-\mu^4 - \mu^3\lambda g_3 + \mu^2\lambda^2 g_2 - \mu\lambda^3 g_1 + \lambda^4 g_0} \ , \\ \lambda_1 &= \frac{-(-\mu^2\lambda - \mu\lambda^2 g_3 + \lambda^3 g_2)}{-\mu^4 - \mu^3\lambda g_3 + \mu^2\lambda^2 g_2 - \mu\lambda^3 g_1 + \lambda^4 g_0} \ , \\ \lambda_2 &= \frac{-(\mu\lambda^2 + \lambda^3 g_3)}{-\mu^4 - \mu^3\lambda g_3 + \mu^2\lambda^2 g_2 - \mu\lambda^3 g_1 + \lambda^4 g_0} \ , \\ \lambda_3 &= \frac{\lambda^3}{-\mu^4 - \mu^3\lambda g_3 + \mu^2\lambda^2 g_2 - \mu\lambda^3 g_1 + \lambda^4 g_0} \ . \end{aligned}$$

We introduce new notation

$$\begin{aligned} g_0 = p_4, \ g_1 = p_3, \ g_2 = p_2, \ g_3 = p_1 \ , \\ G^4 = p_1 G^3 + p_2 G^2 + p_3 G + p_4 \ , \end{aligned} \tag{13.92}$$

then

$$\lambda_0 = \frac{\mu^3 + \mu^2\lambda p_1 - \mu\lambda^2 p_2 + \lambda^3 p_3}{\mu^4 + \mu^3\lambda p_1 - \mu^2\lambda^2 p_2 + \mu\lambda^3 p_3 - \lambda^4 p_4} ,$$
$$\lambda_1 = \frac{-\mu^2\lambda - \mu, \lambda^2 p_1 + \lambda^3 p_2}{\mu^4 + \mu^3\lambda p_1 - \mu^2\lambda^2 p_2 + \mu\lambda^3 p_3 - \lambda^4 p_4} ,$$
$$\lambda_2 = \frac{\mu\lambda^2 + \lambda^3 p_1}{\mu^4 + \mu^3\lambda p_1 - \mu^2\lambda^2 p_2 + \mu\lambda^3 p_3 - \lambda^4 p_4} ,$$
$$\lambda_3 = \frac{-\lambda^3}{\mu^4 + \mu^3\lambda p_1 - \mu^2\lambda^2 p_2 + \mu\lambda^3 p_3 - \lambda^4 p_4} ; \tag{13.93}$$

recall that

$$(\Lambda^{-1})_\rho{}^\alpha = \lambda_0 + \lambda_1 G + \lambda_2 G^2 + \lambda_3 G^3 .$$

Degrees of the matrix G we can associate the following invariants (see Chap. IV in [9]):

$$\begin{aligned}
\mathrm{Sp}(G) = g_1 + g_2 + g_3 + g_4 = s_1 , \quad s_1 = G_\alpha{}^\alpha(x) , \\
\mathrm{Sp}(G^2) = g_1^2 + g_2^2 + g_3^2 + g_4^2 = s_2 , \quad s_2 = G_\alpha{}^\rho(x) G_\rho{}^\alpha(x) , \\
\mathrm{Sp}(G^3) = g_1^3 + g_2^3 + g_3^3 + g_4^3 = s_3 , \quad s_3 = G_\alpha{}^\rho(x) G_\rho{}^\sigma(x) G_\sigma{}^\alpha(x) , \\
\mathrm{Sp}(G^4) = g_1^4 + g_2^4 + g_3^4 + g_4^4 = s_4 , \quad s_4 = G_\alpha{}^\rho(x) G_\rho{}^\delta(x) G_\delta{}^\sigma(x) G_\sigma{}^\alpha(x);
\end{aligned} \tag{13.94}$$

from these place the quantities $g_1, ..., g_4$ stand for four eigenvalues of the matrix G.

Invariants s_i and p_i obey the Newton recurrence formulas [9]:

$$\begin{aligned}
p_1 = s_1 = \mathrm{Sp}(G) , \quad p_2 = \frac{1}{2}(s_2 - p_1 s_1) = \frac{1}{2}\left[\mathrm{Sp}(G^2) - p_1 \mathrm{Sp}(G)\right] , \\
p_3 = \frac{1}{3}(s_3 - p_1 s_2 - p_2 s_1) = \frac{1}{3}\left[\mathrm{Sp}(G^3) - p_1 \mathrm{Sp}(G^2) - p_2 \mathrm{Sp}(G)\right] , \\
p_4 = \frac{1}{4}(s_4 - p_1 s_3 - p_2 s_2 - p_3 s_1) = \frac{1}{4}[\mathrm{Sp}(G^4) - p_1 \mathrm{Sp}(G^3) - p_2 \mathrm{Sp}(G^2) - p_3 \mathrm{Sp}(G)].
\end{aligned}$$

From this it follow the following representations for the invariants p_i:

$$\begin{aligned}
p_1 = \mathrm{Sp}(G) , \quad p_2 = \frac{1}{2}\mathrm{Sp}(G^2) - \frac{1}{2}\mathrm{Sp}^2(G) , \\
p_3 = \frac{1}{3}\left[\mathrm{Sp}(G^3) - \mathrm{Sp}(G)\mathrm{Sp}(G^2) - \frac{1}{2}\left(\mathrm{Sp}(G^2) - \mathrm{Sp}^2(G)\right)\mathrm{Sp}(G)\right] \\
= \frac{1}{3}\mathrm{Sp}(G^3) - \frac{1}{2}\mathrm{Sp}(G^2)\mathrm{Sp}(G) + \frac{1}{6}\mathrm{Sp}^3(G) ,
\end{aligned}$$

$$\begin{aligned}
p_4 = \frac{1}{4}\left[\mathrm{Sp}(G^4) - \mathrm{Sp}(G)\mathrm{Sp}(G^3) - \frac{1}{2}\mathrm{Sp}^2(G^2) + \frac{1}{2}\mathrm{Sp}^2(G)\mathrm{Sp}(G^2)\right. \\
\left. -\frac{1}{3}\mathrm{Sp}(G^3)\mathrm{Sp}(G) + \frac{1}{2}\mathrm{Sp}(G^2)\mathrm{Sp}^2(G) - \frac{1}{6}\mathrm{Sp}^3(G)\mathrm{Sp}(G)\right] ;
\end{aligned}$$

finally for p_4 we find the expression

$$p_4 = \frac{1}{4}\left[\mathrm{Sp}(G^4) - \frac{4}{3}\mathrm{Sp}(G)\mathrm{Sp}(G^3) - \frac{1}{2}\mathrm{Sp}^2(G^2) + \mathrm{Sp}^2(G)\mathrm{Sp}(G^2) - \frac{1}{6}\mathrm{Sp}^4(G)\right] .$$

In the case when the matrix G is antisymmetric, we have the equality

$$\begin{aligned} \tilde{G} = -G,\ p_1 = \mathrm{Sp}G = 0, \\ \tilde{G^3} = -G^3, \qquad \mathrm{Sp}(G^3) = 0\ , \\ p_1 = 0, \qquad p_2 = \frac{1}{2}\mathrm{Sp}(G^2), \\ p_3 = 0,\ p_4 = \frac{1}{4}\mathrm{Sp}(G^4) + \frac{1}{8}\mathrm{Sp}^2(G^2)\ . \end{aligned} \tag{13.95}$$

and the characteristic equation (9.7) takes the form

$$G^4 - p_2 G^2 - p_4 = 0\ , \tag{13.96}$$

this case is realized in the construction of the characteristic polynomial for the electromagnetic tensor. In this case, the relations (13.93) become more simple

$$\begin{aligned} \lambda_0 &= \frac{\mu^3 - \mu\,\lambda^2\,p_2}{\mu^4 - \mu^2\,\lambda^2\,p_2 - \lambda^4\,p_4}\,, \\ \lambda_1 &= \frac{-\mu^2\,\lambda + \lambda^3\,p_2}{\mu^4 - \mu^2\,\lambda^2\,p_2 - \lambda^4\,p_4}\,, \\ \lambda_2 &= \frac{\mu\,\lambda^2}{\mu^4 - \mu^2\,\lambda^2\,p_2 - \lambda^4\,p_4}\,, \\ \lambda_3 &= \frac{-\lambda^3}{\mu^4 - \mu^2\,\lambda^2\,p_2 - \lambda^4\,p_4}\,. \end{aligned} \tag{13.97}$$

For additional verification we consider the simple case in absence of electromagnetic tensor when additionally the space-time is described by the Ricci tensor of the following simple form (elementary examples are the de Sitter spaces)

$$\begin{aligned} G_{\alpha\beta} = \frac{R}{4}g_{\alpha\beta},\ G_\alpha{}^\beta = \frac{R}{4}\delta_\alpha{}^\beta, \\ \mathrm{Sp}G = R, \qquad \mathrm{Sp}(G^2) = \frac{1}{4}R^2, \\ \mathrm{Sp}(G^3) = \frac{1}{4^2}R^3, \qquad \mathrm{Sp}(G^4) = \frac{1}{4^3}R^4, \end{aligned} \tag{13.98}$$

that is

$$\begin{aligned} p_1 = R\,, \quad p_2 = \frac{1}{2}\frac{1}{4}R^2 - \frac{1}{2}R^2 = -\frac{3}{8}R^2\ , \\ p_3 = \frac{1}{3}\frac{1}{16}R^3 - \frac{1}{2}\frac{1}{4}R^2R + \frac{1}{6}R^3 = \frac{1}{16}R^3\ , \\ p_4 = \frac{1}{4}\left[\frac{1}{16\cdot 4}R^4 - \frac{4}{3}R\frac{1}{16}R^3 - \frac{1}{2}\frac{1}{16}R^4 + R^2\frac{1}{4}R^2 - \frac{1}{6}R^4\right] = -\frac{1}{4^4}R^4\ . \end{aligned} \tag{13.99}$$

Expressions for p_i correspond to the following characteristic equation

$$G = (G_\alpha{}^\beta), \qquad (G - \frac{R}{4})^4 = 0 \, . \tag{13.100}$$

Note that in the case of the presence of a geometric background

$$D_\beta \Phi - \lambda \left(F_{\beta\alpha} + i \frac{\hbar c}{e} \frac{R}{4} g_{\beta\alpha} \right) \Phi^\alpha - \mu \Phi_\beta = 0 \, ,$$
$$D^\beta \Phi_\beta - \mu \Phi = 0 \, ; \tag{13.101}$$

and in the absence of an external electromagnetic field, the system of equations (13.101) looks as follows

$$D^\beta \Phi_\beta = Mc\Phi \, , \qquad D_\beta \Phi = \left(Mc + i\lambda \frac{\hbar c}{e} \frac{R}{4} \right) \Phi_\beta \, . \tag{13.102}$$

In particular, in the case of the de Sitter spaces ($R(x) = R$) is equivalent to an effective additive (with a plus or minus) to the mass of the particle

$$D^\beta \Phi_\beta = Mc\Phi \, , \quad D_\beta \Phi = \left(Mc + i\lambda \frac{\hbar c}{e} \frac{R}{4} \right) \Phi_\beta \, ; \tag{13.103}$$

14 Conclusion

Relativistic theory of the Cox's scalar not point-like particle with intrinsic structure is developed on the background of arbitrary curved space-time. It is shown that in the most general form, the extended Proca-like tensor first order system of equations contains non minimal interaction terms through electromagnetic tensor $F_{\alpha\beta}$ and Ricci tensor $R_{\alpha\beta}$.

In relativistic Cox's theory, the limiting procedure to non-relativistic approximation is performed in a special class of curved space-time models. This theory is specified in simple geometrical backgrounds: Euclid's, Lobachevsky's, and Riemann's. Wave equation for the Cox's particle is solved exactly in presence of external uniform magnetic and electric fields in the case of Minkowski space. Non-trivial additional structure of the particle modifies the frequency of a quantum oscillator arising effectively in presence if external magnetic field. Extension of these problems to the case of the hyperbolic Lobachevsky space is examined. In presence of the magnetic field, the quantum problem in radial variable has been solved exactly; the quantum motion in z-direction is described by 1-dimensional Schrödinger-like equation in an effective potential which turns out to be too difficult for analytical treatment. In the presence of electric field, the situation is similar. The same analysis has been performed for spherical Riemann space model.

General conclusion can be done: the effects of large scale structure of the Universe depends greatly on the form of basic equations for elementary particle, any modifications of them lead to new physical phenomena due to non-Euclidean geometry background.

15 Acknowledgements

This work was supported by the Fund for Basic Researches of Belarus, F 13K-079, within the cooperation framework between Belarus and Ukraine. Authors are grateful to organizers of The International conference in honor of Ya. B. Zeldovich 100-th Anniversary: Subatomic particles, Nucleons, Atoms, Universe: Processes and Structure, held March 10–14, 2014, Minsk, Belarus for possibility to give this talk.

References

[1] W. Cox. Higher-rank representations for zero-spin field theories. J. Phys. Math. Gen. 1982. Vol. 15, no 2. P. 627–635.

[2] V.V. Kisel. Exact solutions of the Cox's equation for a particle with spin zero in external electrimagnetic fields. Vesti NANB. ser. fiz.-mat. 200. no 2. P. 82–85 (in Russian)

[3] Red'kov V.M. Field particles in Riemannian space and the Lorentz group. Belarussian Science, Minsk, 2009. 486 pp.

[4] A.A., Red'kov V.M., Krylov G.G.. Schrödinger particle in magnetic and electric fields in Lobachevsky and Riemann spaces. // Nonlinear Phenomena in Complex Systems. 2008. **11**. no 4. 403–416.

[5] V.M. Red'kov, E.M. Ovsiyuk. Quantum mechanics in spaces of constant curvature. Nova Science Publishers. Inc., New York, 2012. - 434 pages.

[6] E.M. Ovsiyuk, O.V. Veko. On behavior of quantum particles in the electric field in spaces of constant curvature, hyperbolic and spherical models. Ukr. Phys. J. 2013. **58**, no 11. 1065–1072.

[7] H. Bateman, A. Erdélyi. Higher transcendantal functions. Vol. I. – Russian translation, Moscow, 1965.

[8] H. Bateman, A. Erdélyi. Higher transcendantal functions. Vol. II. – Russian translation, Moscow, 1966.

[9] F.R. Gantmakher. Theory of Matrices. – 4th ed. – .: Nauka, 1988. – 552 pp.